

# A Review of Bisphenol A Effects on White Adipose Tissue and Its Correlation with Various Cancers

Amir Karbalaee Hasani , Zivar Nejad Ebrahimi , Neda Mahami , Mohammad Alizadeh ,  
Zahra Rasooli , Mina Hemmati 

Dept. of Biochemistry, School of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran

## Article Info

 [10.30699/jambs.30.139.86](https://doi.org/10.30699/jambs.30.139.86)

Received: 2020/11/07;

Accepted: 2021/03/15;

Published Online: 31 Jan 2022;

Use your device to scan and read the article online



## Corresponding Information:

**Mina Hemmati,**

Dept. of Biochemistry, School of Medicine, Zanjan University of Medical Sciences, Zanjan, Iran

E-Mail: [mina1hemmati@yahoo.com](mailto:mina1hemmati@yahoo.com),  
[mhemmati@zums.ac.ir](mailto:mhemmati@zums.ac.ir)



Copyright © 2021. This is an original open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permits copy and redistribution of the material just in noncommercial usages with proper citation.

## Introduction

Many synthetic chemicals are marketed aimed at improving the living standards without considering the negative consequences on human health (1). Bisphenol A (BPA) (4, 4'-dihydroxy-2, 2 di phenyl propane), as a synthesized chemical (material) is formed by the reaction of two phenol groups with an acetone molecule (2, 3). BPA is an estrogen monomer with a molecular weight of 228.29 g / mol and is known by the chemical formula of C<sub>15</sub>H<sub>16</sub>O<sub>2</sub>. BPA is a white crystalline substance with a mild phenolic odor and a melting temperature of 153-159 °C (4, 5) which is used as a primary substance for the production of the polycarbonate plastics, epoxy resins, and polyester. Polycarbonates are used in making water bottles, baby food bottles, toys, heat plates, home appliances, sanitary ware, and medical equipment (5). BPA is also utilized in the production of various plastics, such as polyvinyl chloride, which is used in the manufacturing of medical pipes, water pipes, and as a lining in the dentistry (5). Exposure to BPA is mainly through the hydrolysis of plastic containers containing the polycarbonate and epoxy resins. BPA monomers obtained

## ABSTRACT

Bisphenol A, as a primary substance used for making many plastic devices enters the body through various routes such as skin penetration, oral ingestion, and inhalation and is among the chemicals leading to the disruption of the endocrine system. It exerts its destructive effects on the various tissues through estrogen, androgen, thyroid, pregnane X, and aryl hydrocarbon receptors. This review study was conducted to evaluate the pharmacological effects of bisphenol and its signaling pathways especially in the adipose tissue. Studies have shown that the disruption in the level of adipocytokines can contribute to the formation and progression of the chronic diseases such as cancers. Decreased level of adiponectin (as an adipokine secreted from white adipose tissue) and increased levels of IL-6, TNF- $\alpha$ , and leptin are among the disruptions caused by bisphenol which can develop different types of cancers. Accordingly, it seems that the use of bisphenol and contact with it should be reconsidered with regard to its destructive effects on the endocrine system.

**Keywords:** Bisphenol A, Adipocytokines, Adipose tissue, Cancer


from BPA-produced polymers can be combined with the foods and water; and the consumption of these contaminated foods is the first source of human exposure to BPA (6). Absorption of BPA by the human has been estimated by 10  $\mu\text{g} / \text{kg} / \text{day}$  (7). During 1995 - 2014, BPA production in the United States and Europe has increased by 37 and 84%, respectively, and this process is expected to increasingly continue (8). Given the human contact with BPA and its harmful effects on the human health, the Environmental Protection Agency (EPA) has introduced BPA as a destructive chemical against the humans' endocrine system (9).

### 1. Pharmacokinetic of the BPA

In the United States, BPA has been observed in the urine samples of more than 92% of the people at 6 years of age and older (10, 11). The US EPA has stated the reference dose of 50  $\mu\text{g} / \text{kg} / \text{day}$  for BPA. Meanwhile, the European Food Safety Authority (EFSA) has reduced the safe exposure rate to BPA from 50 to 4  $\mu\text{g} / \text{kg} / \text{day}$  (11). Following ingestion, about 90% of BPA passes through

the liver and its primary metabolism occurs in the intestine and liver (12,13). Glucuronidated and sulfonated BPAs are among the major secondary metabolites of BPA which are excreted through the urine (Figure 1). BPA mostly acts in the humans' body through binding to the steroid hormone receptors, such as estrogen and androgen receptors to alter the gene expression. The distribution of

these receptors varies in different tissues. Therefore, the binding of BPA to these receptors creates a variety of responses on the cell surface. BPA can bind to these receptors both freely and in the conjugated form in the bloodstream and then, exert its physiological functions (13,14).



**Figure 1.** Chemical structure of BPA and its primary human metabolites, BPA-glucuronide. In humans, elimination of BPA is via urine in the form of BPA or BPA conjugates, mostly BPA-glucuronide. The conjugation is done by UDP-glucuronyl transferase (UGT) and sulfotransferase (SULT). Figure was prepared using templates on the servier medical art website (<http://smart.servier.com/image-set-download/>).


## 2. BPA as an Endocrine Disrupting Compound

Endocrine Disrupting Compounds (EDCs) are the compounds altering the function of the hemostatic and hormonal systems. EDCs interfere with the function of the nuclear receptors, estrogenic non-nuclear receptors, non-estrogenic receptors, and neurotransmitter receptors, such as serotonin, dopamine, and norepinephrine receptors, and also the orphan receptors including aryl hydrocarbon receptor as well as the pathways involved in the steroid synthesis or metabolism and other mechanisms converging with the endocrine and reproductive systems (15).

### 2.1. Estrogen Receptors

BPA can exert its physiological effects through various physiological receptors including the Estrogen Receptors alpha (ER $\alpha$ ), Estrogen Receptors beta (ER $\beta$ ), Estrogen-Related Receptors gamma (ERR  $\gamma$ ), and Peroxisome Proliferator-Activated Receptor gamma (PPAR  $\gamma$ ) (16). Compared to 17 $\beta$ -estradiol (E2), BPA tendency for

binding to ER $\beta$  is higher than ER $\alpha$  (17). BPA acts as an ER $\alpha$  agonist and creates the same change as E2, while preventing the formation of the correct conformation of the ligand-binding domain in ER $\beta$  and acting as an ER $\beta$  antagonist (18, 19). Given the effect of E2 on induction of the cellular proliferation, the effect of BPA has also been investigated in this field. In fact, the balance between signaling of ER $\alpha$  and ER $\beta$  and the level of these receptors causes the balance between the cell death and proliferation regulated by E2 (20). BPA acts as a proliferative agent by mimicking the E2 function in the presence of ER. While in the presence of ER $\beta$ , BPA acts as an antagonist of the E2-ER $\beta$  complex indicating that BPA leads to the increase in the cell growth through ER $\alpha$  and on the other hand, it reduces the cell death through the antagonistic activity during binding to ER $\beta$  (Figure 2). These two functions of BPA in the presence of ER $\beta$  and ER $\alpha$  can create changes in the cells resulting in the development of cancer (21).



**Figure 2.** Interaction of BPA with ER $\alpha$ /ER $\beta$  resulting in the development of various cancers. In the presence of ER $\alpha$ /ER $\beta$  BPA binds to these receptors and induces cell proliferation.

## 2.2. Androgen Receptor (AR)

BPA acts as an androgen receptor antagonist. Evidence indicates that the antagonistic activity of BPA on the androgen receptor prevents the transmission of the receptor dimer to the nucleus and reduces the expression of the synaptic proteins [Synapsin1 and Postsynaptic Density Protein 95 (PSD95)] in the hippocampus of the male rats resulting in a defect in the spatial memory (22). There is insufficient information on the effect of BPA on the AR transcription activity, and there is not sufficient knowledge about the ability of these compounds to interfere with the androgen-dependent extra-nuclear signals (23). Androgen signaling appears to be less influenced by the intervention of BPA, but BPA can intervene in the treatment of the patients with advanced prostate cancer through mutated ARs (24).

## 2.3. Thyroid Receptors

BPA has a low affinity for binding to the thyroid receptors (TR) and reduces the self-induction effect of the TR through antagonistic activity and at the same time increases the serum thyroxine (22). BPA inhibits the stimulated transcription activity by Triiodothyronine (T3) through antagonistic activity. Through a Nuclear receptor Co repressor (NCoR), BPA could reduce the expression of the TR gene (5). According to the literature, BPA can influence the expression of the genes involved in the brain development by interfering with the thyroid hormone receptor (25). These effects are mediated by the receptors located in the nucleus, where they act as transcription suppressors in connection with DNA through Silencing Mediator of Retinoic acid and Thyroid hormone receptor (SMRT) and NCoR. The extra-nuclear activity of the TR depends on integrin  $\alpha\beta_3$ , which is responsible for phosphorylation and activation of the nuclear TR through activating the Mitogen-Activated Protein Kinase (MAPK) / cellular Sarcoma (c-Src) pathway (26). BPA directly interferes with the  $\beta_3$  / integrin / c-Src / MAPK / TR- $\beta_1$  pathway at low concentrations, which could affect TR signaling.

## 2.4. Pregnane X Receptor

Human Steroid and Xenobiotic Receptor / rodent Pregnane X Receptor (SXR / PXR) and Constitutive

Androstane Receptor (CAR) are the two important regulators for metabolism of the xenobiotic and steroid hormones (19). These receptors in the liver and intestine mediate the induction of the cytochrome enzymes p450 and transporters such as P-glycoprotein in response to xenobiotic ligands and steroid hormones. SXR / PXR are unusual nuclear receptors, which have extensive ligand specificity and are activated by a lot of EDCs such as BPA. Activation of the SXR / PXR and CAR, and consequently an increase in the expression of their target genes by various components causes an increase in the levels of EDCs and at the same time changes the rate of access to estrogens and androgens (27). In fact, BPA acts as a strong agonist for hPXR. Combination of the effect of BPA and other EDCs results in the synergistic effects on the hPXR function (28).

## 2.5. Aryl Hydrocarbon Receptor

Aryl hydrocarbon receptor binds to several chemicals and indirectly influences the metabolism of xenobiotics as well as the synthesis and metabolism of steroids. Aryl hydrocarbon receptors are the transcription factors mediating the effects of various chemicals most importantly, Tetrachlorodibenzodioxin. After binding the ligand to the AHR in the cytoplasmic segment, it enters the nucleus by the corresponding transmitter (ARNT) which binds to the responsive elements in DNA and leads to the transcription of the genes involved in the xenobiotic metabolism including CYPs. The AHR repressor must be bound to ARNT to disable this process or the messaging path. BPA disrupts the AHR expression and its activity in the embryo via AHR upstream transcriptional regulation (29). BPA results in the changes in the primary adipogenesis along with hypertrophic modification of the adipocytes and overexpression of the lipogenic genes such as PPAR  $\gamma$ , Sterol Receptor Element-Binding Protein 1C (SREBP-1C), Lipoprotein Lipase (LPL), and Fatty Acid Synthetase (FAS) (30). BPA is able to increase the expression of the PPAR  $\gamma$  -related-mRNA in the 3T3-L1 pre-adipocytes (31). The cAMP Response Element-Binding protein (CREB) also plays an important role in the differentiation of the adipocytes, and the effects of BPA on the differentiation of the adipocytes are likely to occur through the rapid activation of CREB. Exposure to low BPA concentrations for a long time influences the

adipogenic differentiation program by increasing the peri-adipocyte growth and altering the genes regulating the adipogenesis (25). ERR $\gamma$  is present in the adipose tissue. Evidence indicates that ERR $\gamma$  with its two isoforms (ERR $\alpha$  /  $\beta$ ) plays an important role in controlling the energy homeostasis and biogenesis and the natural function of the mitochondria. Studies have shown that BPA exerts its effects on the body weight regulation through ERR $\gamma$  (30). BPA binds strongly to the human ERR, and it also binds to ERR $\gamma$  at low concentrations while maintaining the ERR  $\gamma$  activity in the presence of 4-hydroxydebrisoquine (4-OHD). The effects of BPA at low concentrations on the energy homeostasis are likely mediated by these nuclear receptors (19). BPA acts in the body mainly through the estrogen receptor. This receptor is expressed in various tissues of the body. BPA can cause the disruptions in the tissues of these estrogen receptors by binding to these receptors (32).

### 3. Expression of the ERs in the Adipose Tissue

Hormones are the main regulators of the adipose tissue function and play an important role in the evolution and development of the adipocytes. A wide range of growth hormones and factors, such as thyroid hormone, glucocorticoids, catecholamines, insulin, glucagon, and insulin-like growth factors modify the activity and development of the adipose tissue. Estrogens, especially E2 play an important regulatory role in the metabolism and distribution of the adipose tissue (33). The sex steroids contribute to the function of the white adipose tissue through controlling the lipolysis, lipogenesis, adipocyte differentiation, insulin sensitivity, and production and secretion of the adipokines (34). In adults, increased estrogen plasma concentration is associated with the decreased food intake and body weight, and estrogen deficiency is accompanied by the initiation of the menopause in overweight women (35). Studies on the humans and laboratory animals have suggested that estrogen is the most important regulator of the white adipose tissue in women (36). Estrogens exert their effects on the adipose tissue through the steroid receptors (37). ER $\alpha$  was reported as the first receptor for estrogen in 1973. The second steroid receptor, ER $\beta$  was identified in 1996. Studies on the laboratory animals such as adult rats have shown that the endogenous E2 is the most abundant and predominant estrogen, and exerts its inhibitory effect on the number of adipocytes and lipogenic cells through the ER $\alpha$  receptor (35). Both ER $\alpha$ / $\beta$  receptors are expressed at the mRNA and protein levels in the human adipose tissue with ER $\alpha$  predominance. There is no regional or gender difference in the adipose tissue for ER $\alpha$  expression, while ER $\beta$  expression is higher in women than men (38). Different roles have been attributed to ER $\alpha$  and ER $\beta$  regarding exerting the estrogen effects on the adipose tissue. Obesity has been shown to increase in therats without ER $\alpha$ , while normal adipose tissue mass was observed in the rats without ER $\beta$  (39). In the studies on the 3T3-L1 cell line, ER receptors have also been observed in the membrane and cell nucleus (40). Estrogens can influence the adipose tissue activity in several ways:

1. A genomic response characterized by the changes in the gene transcription that occurs from several hours to several days.

2. Non-genomic signal events that occur from several seconds to several minutes after cell stimulation (21).

With regard to the presence of the ERs in the adipose tissue and their activities in the adipose tissue, BPA can cause the disruptions in the adipose tissue by the ERs. These disruptions include the changes in the normal secretions of adipocytokines from this tissue (41).

### 4. Disruption in the Secretions of the Adipose Tissue Leading to the Development of Cancer

The adipose tissue is a dynamic tissue involved in regulation of the glucose and lipid metabolism, energy homeostasis, and inflammation. In addition, the adipose tissue is an active and complex endocrine organ playing a role in the metabolism of sex steroids and the synthesis and secretion of the active adipokines (42). In addition to storing fat, the white adipose tissue is an active endocrine gland that secretes more than 50 cytokines and other molecules generally called adipocytokines including adiponectin (APN), leptin, Tumor Necrosis Factor  $\alpha$  (TNF $\alpha$ ), and Interleukin 6 (IL-6) (43, 44). There is a special relationship between the adipose tissue and different types of cancer including colorectal cancer, kidney cancer, postmenopausal breast cancer, esophageal cancer, etc. (45). The balance between the number of adipocytes and cancer cells is created by cytokines including IL-6, TNF $\alpha$ , and adipokines, such as APN and other proteins produced from the adipose tissue to control invasion and proliferation of the cancer cells (46). Recent studies have confirmed the effects of the adipocytokines in cancer (47). According to the recent studies, BPA can cause changes in the secretions of the adipose tissue through different cellular pathways (1). The following investigates the effects of BPA on adipocytokines and the role of these changes in cancer in detail.

### 5. The Effects of BPA on Cytokine Secretion

Numerous reports have shown that BPA influences the levels of important hormones associated with metabolism, such as APN and leptin which are among adipocytokines (48). EDCs can disrupt the signaling by the regulatory hormones including APN and leptin (32). Regarding the presence of BPA around the world as well as the constant exposure of humans to the it, inhibition of APN secretion, and increased levels of the IL-6, TNF $\alpha$ , and leptin at nanomolar concentrations may be the important factors in the endocrine disruption and homeostasis of metabolism (18).

#### 5.1. The Effects of BPA on the APN Secretion from the Adipose Tissue

APN is the most abundant adipocytokine secreted by the adipocyte cells of the adipose tissue. APN, as an adipokine is responsible for energy storage in the form of TG, and has anti-apoptotic and anti-inflammatory activities (49). Clinical studies have shown that the APN level plays an important role in the protection against the

carcinogenic factors. Some cancer cells influence the APN receptors and their downstream signaling pathways to prevent the apoptosis (50). APN can prevent the development of the cancers such as breast cancer, lung cancer, thyroid cancer, and colorectal cancer by regulating the signaling pathways, apoptosis, and the cell cycle (51). Atrial Natriuretic Peptide (ANP) is a mediator between the AMP-activated Protein Kinase (AMPK), Janus Kinase (JAK) / Signal Transducer and Activator of Transcription 3 (STAT3), and MAPK signaling pathways (51, 52). ANP causes the activation of apoptosis and inhibition of the cell proliferation through the AMPK signaling pathways (52). Studies conducted on humans and animals have shown that estrogen reduces APN secretion. Results of a study on the rats without the aromatase enzyme showed that the APN levels were reduced by E2 consumption (53). EDCs act directly on the steroid hormone receptors or indirectly through non-steroidal receptors. Like E2, BPA activates the ER $\alpha$ -dependent signaling pathway leading to the MAPK / Extracellular-Regulated Kinase (ERK), and Protein Kinase B (AKT) phosphorylation (54). Low BPA concentrations inhibit the APN secretion in the human adipose cells *in vitro*. It has been observed that BPA accumulates in the adipose tissue. In addition, the ERR $\beta$  receptor, as a nuclear receptor has a high tendency for binding to BPA (55).

BPA regulates the expression and phosphorylation of AKT acting on the downstream of the Phosphoinositide 3-Kinase (PI3K), and the PI3K signaling pathway is accompanied by the production and secretion of APN. The PI3K-AKT pathway is activated when the insulin hormone binds to its receptor at the cell level that plays a key role in regulating the gene expression and APN secretion. In fact, BPA inhibits the expression and phosphorylation of AKT, and APN levels have been reported to be significantly lower in the cell cultures treated with BPA. BPA does not act through the nuclear ER pathway in the 3T3-L1 cells, but it also binds to the estrogen-related receptor in the human (56). In fact, PI3K, MAPK, and ER have been reported as the main components of the signal transmitter for E2 or as foreign estrogens in the regulation of the genes in the 3T3-L1 cell line (40). Both BPA and E2 inhibit the APN secretion, but they do not act through a single mechanism (32). APN synthesis is stimulated by PPAR $\alpha$ , insulin, and Insulin-like Growth Factor-1 (IGF-1) agonists. Increased expression of APN by PPAR $\alpha$  is highly regarded, because being an antagonist for PPAR $\alpha$  is among the mechanisms through which BPA can suppress the APN secretion. The APN secretion is controlled at three levels: biosynthesis, assembly, and release (57). The APN synthesis is stimulated by PPAR $\alpha$ , insulin, and IGF-1 agonists and is inhibited by TNF- $\alpha$  and catecholamines (18). Increase in the APN expression is important because the mechanism by which BPA can suppress the APN secretion is performed by the PPAR $\alpha$  antagonist. In fact, conjugated compounds with BPA are potential antagonists for PPAR $\alpha$  activities in the adipocytes (58). In addition, BPA can inhibit the Protein Disulfide Isomerase (PDI) enzyme

playing an important role in the formation of the disulfide bonds in APN, which is considered as a mechanism involved in the reduction of APN by BPA (59).

### 5.1.1. Decreased Level of APN and Its Association with Various Cancers

#### Endometrial Cancer (EC)

There is an inverse relationship between the serum level of APN and the risk of endometrial cancer (60). APN reduces the viability of the EC cells and decreases their number through inhibiting the proliferation and inducing the apoptosis (50). Previous studies have shown an inverse relationship between the serum APN level and the risk of EC in the women under 65 years of age (51, 61, 62).

#### Hepatic Cancer (HC)

The risk of liver cancer is directly associated with the serum level of APN (50, 63). APN responds to liver cancer by preventing the cell proliferation and pro-apoptotic factors. The expression of Cyclin D1 and Proliferating Cell Nuclear Antigen (PCNA) decreases and instead, the expression of Caspase-3 increases in the liver cancer cells. APN influences the hepatocellular cancer cells through AMPK, c-Jun N-terminal Kinase (JNK), and mammalian Target of Rapamycin (mTOR). APN causes the JNK phosphorylation in the HepG2 and Huh7 cells and increases the AMPK activity through phosphorylation (63).

#### Prostate Cancer (PC)

Prostate cancer is one of the most common cancers in men metastasizing to other organs. Clipping the androgen pathway is among the therapeutic methods for treatment of PC; however, this therapeutic method does not respond in some cases. APN level is lower than the normal in the patients with PC and is inversely related to the disease progression (50, 61, 63-65). The mRNA expression of AdipoR1 / R2 receptors is also reduced in the PC cells. Serum APN level and profile amount of its receptors can predict the risk of PC (51, 52, 65). APN influences the pathogenesis of PC through the JNK and STAT3 pathways. ANP causes the activation of the JNK, AMPK, and PI3 kinase / AKT pathways in the PC cells. The AMPK pathway inhibits the growth and proliferation of the PC cells through 5-aminoimidazole-4-carboximid ribose (AICAR) (51, 63, 66).

#### Gastric Cancer (GC)

Low serum level of APN is directly associated with an increased risk of gastric cancer. Increased expression of the AdipoR1 receptor suppresses the growth of the GC cells (50, 51). Serum level of APN is lower than the normal range in the patients with esophageal cancer. Esophageal cancer is among the malignant cancers metastasizing to the gastrointestinal tract and in most cases, the patients die within 5 years (50, 67). According to literature, low serum level of APN also increase the risk of pancreatic cancer (50).

#### Colorectal Cancer (CRC)

Numerous studies have proved an inverse relationship between the serum level of APN and the risk of colorectal cancer (45, 46, 50, 63, 66, 68, 69). CRC is among the most common cancers associated with obesity. Previous studies have shown that low serum level of APN is strongly associated with an increased risk of CRC (70). AdipoR1 / R2 are expressed in the CRC cells and also involve the lymph nodes. In the low stages of the disease, the rate of AdipoR1 expression is associated with the tumor size (46, 63, 64). High serum level of APN reduces the risk of CRC by 60% (50, 71). Inhibition of the Cyclin-Dependent Kinase (CDK) inhibitors, P21 / P27 through AdipoR1 / R2 increases the cell cycle suppression. Therefore, APN can control the proliferation of the cancer cells by activating several tumor suppressor genes (68).

#### Thyroid Cancer (TC)

Thyroid cancer is the second most common cancer in young women. APN receptors are expressed both in the healthy thyroid and TC cells (46). Plasma APN level is lower in patients with TC than the healthy individuals (60).

#### Renal Cell Cancer (RCC)


Low serum level of APN not only increases the risk of kidney cancer but also causes metastasis. The expression of AdipoR1 / R2 receptors is lower in the kidney tissues than the other tissues resulting in a decrease in the protective potential of APN in kidneys (50, 60, 64, 70, 72).

### 5.2. The Effects of BPA on the Leptin Secretion from the Adipose Tissue

Leptin, as a 16 -kDa peptide hormone contains 167 amino acids. The three-dimensional structure of leptin consists of a motif containing four alpha helices, which is very similar to the structure of the (IL-6) family (73). In terms of the structure, leptin is a typical neuropeptide with anorexic function. It is one of the important molecules involved in the development of obesity, which is also called the "satiety hormone" because it plays a key role in absorbing the nutrients and controlling the energy consumption. It is derived from the Greek word "leptos", which means the slender and thin (74). Vokberg et al., showed that the exposure to BPA leads to weight gain and accumulation of the adipose tissue mass, it also creates some changes in the leptin level (75). Increased serum level of leptin is associated with several cancers including CRC, PC, and Breast Cancer (BC) (76, 77). Leptin increases the proliferation, migration, and invasion of the cancer cells (69, 78) and is able to induce the vascularization, increase the level of the Vascular Endothelial Growth Factor (VEGF), and suppress the secretion of the anti-inflammatory cytokines (79). Binding the leptin to its receptor activates several signaling pathways in the form of a cascade. So that, first, tyrosine kinase is activated, and influences the cytokine receptor and binds it to the transcription activators of STAT. STAT migrates toward the nucleus and induces the expression of the genes, such

as down-regulators (including the cytokine signaling inhibitors) like JAK2-STAT3, MAPK, and PI3K / AKT (80). Studies have shown that the leptin levels in the bloodstream are higher in women than men. An increase in the mRNA of Ob (human Obesity) gene in the presence of E2 reflects the direct effect of estrogen on the transcription of Ob gene. Evidence indicates that there are several responsive elements to the Specificity Protein 1(Sp1) in Ob gene promoter in humans. Thus, ER-Sp1 relationship represents the transcription by estrogen and without the need for direct binding of ER to the DNA. In -vitro studies have shown that the treatment of adipocyte cells of the rats with estrogen increases the leptin secretion (81). Specific activities of leptin in target tissues are conducted by its interaction with specific receptors, leading to the secondary activation of the JAK / STAT, MAPK / ERK, and PI3K / AKT pathways (82).

Previous studies have shown that BPA and leptin can initiate similar intracellular messaging pathways that induce cell proliferation. It has also been shown that BPA increases the leptin receptor expression and, like leptin, induces proliferation by activating the STAT3, ERK1 / 2, and AKT messaging pathways. Similar to one of the xenoestrogens, BPA increases the expression of the leptin receptor gene in the ovarian cancer cells and activates the intracellular signaling pathways (83, 84). There is normal overexpression in the amount of leptin and its receptor in the advanced stages of the BC (83). Leptin can bind to all its four receptor forms (ob-Rd, ob-Rc, ob-Ra, ob-Rf) and activate the JAK pathways, and phosphorylate the Insulin Receptor Substrates (IRS), and initiate the AKT / PI3K phosphorylation. Increased expression of the leptin receptor in the cancer cells is associated with the decreased survival (69). Leptin inactivates various signaling pathways including the AKT / PI3K, JAK / STAT, and MAPK / ERK pathways (76, 83-85). The AKT / PI3K pathway plays an important role in regulating the cell growth and preventing the cell proliferation through apoptosis (76). Estrogen is among the compounds that can influence this pathway, which prevents the apoptosis by activating the PI3K / AKT and MEK / ERK pathways, as a common pathway between leptin and estrogen (78). Leptin increases the D1 and c-Myc Cyclin expression, and also increases the rate of cell cycle progression by reducing G1. Cyclin D1 is overexpressed in the BC cells. Cyclin D1 plays an essential and important role in regulating and advancing the cell cycle from stage G1 to S (78). Leptin reduces apoptosis through upregulation of the growth stimulants such as the Cyclin D1 (to intensify the cell proliferation) and downregulation of the tumor suppressors such as P53 and P21 (78). Leptin produced by the adipose tissue adjacent to the tumor causes a local increase in the leptin level and stimulates the tumor growth. In other words, the secretions of the adipose tissue around the tumor into the intercellular space play an important role in the local growth of the tumor (69). The effects of BPA on adipokine secretion and related signaling pathways are summarized in [Figure 3](#).



**Figure 3.** Schematic representation of estrogenic activity of BPA in adipocytes through genomic and non-genomic pathways, as described in the text. BPA mediates transactivation of OB gene. BPA also inhibits APN expression through inhibition of expression and phosphorylation of AKT. Figure was prepared using templates on the servier medical art website (<http://smart.servier.com/image-set-download/>).

### 5.2.1. Increased Level of Leptin and Its Association with Various Cancers

#### Colorectal Cancer (CRC)

Leptin is usually released within the cycle of 2 - 3 hours after a meal, and serum leptin is directly related to the fat reserves; that is, it increases under the obesity conditions and decreases following the weight loss. It has been observed that if the number and size of the fat cells increase, the LEP gene is activated with the synthesis of leptin, which is then secreted into the bloodstream. CRC occurs more in the obese people than the healthy people. Numerous reports have shown that leptin plays an important role in the progression and pathogenesis of CRC. Leptin receptor expression has been observed in the cytoplasm and the cell membrane of many tumor cells including the CRC cells. Thus, it can be said that, the leptin receptors are expressed in 77-95.5% of CRC cases (84). Evidence indicates that obesity has a negative effect on the cancer treatment and leads to the development of drug-resistance, especially through leptin and its receptor. Leptin adipokine, has been suggested to strengthen the tumor and therapeutic resistance in many cancers. Cytokine leptin increases the pre-inflammatory and immune secretion of the cancer cells (86). Leptin also

forms a tumor microenvironment to strengthen the angiogenesis and metastasis. In addition, leptin is associated with the expansion of the cancerous stem cells population, and the expression of the leptin receptor is essential for maintaining the properties of its cancerous stem cells so that, leptin production increases the soluble Intercellular Adhesion Molecule-1 (sICAM-1) in the cancerous cells. ICAM-1 expression has not been observed in the lung and breast cancerous cells, and this effect is regulated by the leptin receptor (ObR). In addition, leptin prescription causes an increase in the expression of ERK, JAK1 / 2 STAT3, and Focal Adhesion Kinase (FAK) genes and also leads to the incidence of ICAM-1 expression (86). Numerous studies have confirmed that the Leptin Receptor (LEPR) signaling can cause the adhesions, angiogenesis, migration, and survival of the cancerous cells. A significant increase in the LEPR expression in patients with metastatic forms of CRC indicates that the LEPR activity is a valuable diagnostic indicator of the metastatic potential of CRC (74).

#### Breast Cancer (BC)

Leptin is the most important mediator of obesity and breast cancer, leading to tumorigenesis, development, growth, and metastasis. Molecular mechanisms assoc-

iated with obesity and BC are complex mechanisms, but four mechanisms can explain this relationship including the expression of the leptin receptor, chronic inflammation of the adipose tissue, sex hormone replacement (IGF-I), and insulin signaling. Leptin increases the risk of BC by cross-reacting with other signaling molecules including ER $\alpha$ , growth and Notch factors, and inflammatory factors. Cross signaling between leptin and IGF-I causes invasion of BC through the reaction of the Epidermal Growth Factor Receptor (EGFR). Mutual signaling between the leptin-Notch and interleukin-1 (NILCO) causes the development of the cancer. Leptin – NILCO axis is a mediator of activation of the stem cells. Signaling of the cancer stem cells strengthens the expression of the leptin receptors by activating the STAT3 pathway and increases the expression of the stem cells' markers such as the octamer-binding transcription factor 4 (OCT4) and (sex determining region Y)-box 2 (SOX2) (87).

### Ovarian Cancer

Leptin and its receptor are overexpressed in the epithelial ovarian cancer, which is significantly associated with the progress-free survival. Increased expression of the leptin receptor gene and protein in the ovarian cancer cells occurs at nanomolar concentrations. Previous studies have shown that leptin and BPA inhibit the expression of the Caspase -3 gene in the OVCAR-3, as the ovarian carcinoma cell line. However, there is no information on the interaction between BPA and leptin in inhibiting the apoptosis in cancer. Activation of the JAK / STAT, MAPK / ERK1 / 2 ,and PI3K / Akt signaling pathways has been investigated previously, and it has been hypothesized that two distinct combinations of leptin and BPA (obesity-related) may regulate the apoptosis through activation of the intracellular targets in the OvCAR-3 ,as the ovarian cancer cells. It has been shown that BPA and leptin interact to inhibit the gene and protein Caspase -3 through different intracellular signaling pathways so that leptin works through STAT-3, while BPA works through ERK1 / 2 (83).

### Thyroid Cancer (TC)

Leptin may influence the migration of the thyroid cells. Also, adipokines, such as APN, leptin, and hepatocyte growth factors are able to regulate the proliferation and invasion of the cancerous cells and may play a role in the development of TC. Leptin serum level increases in the patients with Papillary Thyroid Cancer (PTC). Decreased APN level known as the angiogenesis inhibitor is accompanied by the progression of TC under obesity conditions, and changes in the thyroid function and potential mediators, such as cytokines, insulin, leptin, and APNs may be related to an inflammatory condition (79). Numerous clinical studies have shown that the lymphatic invasion is more frequently observed in the cancers with high leptin expression in different types of tumor. Growing evidence shows that leptin increases the invasion and proliferation in the breast, ovary, lung, bile, and pancreatic cancers and other cancers. Previous studies have demonstrated that leptin is expressed in the cell categories

prone to the migration, and exogenous leptin also increases the ability of the glioma cells to migrate. In addition, it has been reported that leptin plays a role in the activity of the cancerous fibroblasts and the expression of the matrix metalloproteinases greatly influencing the metastatic behavior of the cancerous cells. In addition, there is also evidence showing that leptin increases the transmission of the epithelial cells through various signaling mechanisms leading to the tumor invasion and metastasis (86).

### 5.3. The Effects of BPA on the IL-6 Secretion from the Adipose Tissue

IL-6, as a 184-aa cytokine with a molecular weight of about 25 KDa and pleiotropic function has both anti-inflammatory and pre-inflammatory properties in the immune and non-immune cells (88, 89). IL-6 is synthesized in the immune cells and the peri-adipocytes and adipocytes. IL-6 production is mediated through stimulation by insulin through the cyclic Guanosine Monophosphate (cGMP) / Protein Kinase G (PKG) / CREB pathway, also requiring MAPK (86). Evidence indicates that nanomolar concentrations of BPA may induce a similar response of inflammation in the human adipose tissue. Exposure to BPA causes the secretion of the inflammatory factor of IL6. The activation of normal inflammatory pathways is another finding associated with the disruptions caused by BPA. In fact, the JNK, JAK / STAT, and (nuclear factor kappa light chain enhancer of activated B cells) NF-KB pathways have been shown to be activated in the adipose tissue treated with BPA (86). It is assumed that BPA can stimulate the Toll-Like Receptors (TLRs) and leads to the proper regulation of pre-inflammatory factors such as IL-6 through downstream signaling of JNK and NF-KB (86). BPA may activate the intracellular signaling pathways through TLR and ER ,or GPRC30, leading to the upregulation of IL6, which in turn can stimulate and activate the common inflammatory pathways such as JNK and JAK / STAT pathways (86). After binding of IL-6 to its receptor (IL-6R), it is bound to another receptor called glycoprotein 130 (gp130) and then, signaling is started (86). Dimer form of gp130 activates the tyrosine kinase of JAK1 leading to the activation of the other signaling pathways including the STAT1, PI3K, MAPK, and STAT3 pathways (88-90). STAT3 acts as an oncogene and is the linker between inflammation and cancer. Activation of SATA3 by IL-6 increases the expression of Cyclins B1, D2, D1 ,and C-Myc and inhibits the expression of the CDKs, as a result of which the cell enters the cell cycle. It also increases the expression of a number of survival proteins including the (B-cell lymphoma 2) Bcl-2, (myeloid leukemia cell differentiation protein) Mcl-1, Survivin, and (B-cell lymphoma-extra-large) Bcl-xl. Increased expression of these proteins and continuation of the STAT3 activity lead to the drug resistance in cancer patients. STAT3 activity induces the expression of the Hypoxia-Inducible Factor  $\alpha$  (HIF $\alpha$ ) dependent on the VEGF downstream protein of this pathway (88, 91). Activating the JAK-STAT3 signaling pathway leads to an increase in the expression of the matrix metalloproteinase (MMP7, MMP2, and



MMP9). MMP9 plays an important role in the destruction of the extracellular matrix and causes invasion of the tumor cells (91). Cancer is also an inflammatory disease, and the expression of the cytokines plays a major role in the incidence of cancer (88, 92). Increased expression of IL-6 is associated with the cancers such as colon, lung, prostate, liver, etc (88).

Twenty percent of the cancers are caused by the chronic inflammation. IL-6, as one of the pro-tumorigenic cytokines controls and regulates the proliferation of the cancer cells by regulating the signaling pathways of growth factors including EGF and HGF family members (91). TNF $\alpha$  increases the IL-6 expression during the inflammatory response (88). Increased expression of IL-6 in several organs such as the brain, lungs, the liver, and the bone marrow causes absorption and metastasis of the circulating tumor cells in these organs. Activation of NF- $\kappa$ B in the Kupffer cells stimulates the production of IL-6 causing the metastasis of the lung cancerous cells to the liver (91).

### 5.3.1. Increased Level of IL-6 and Its Association with Various Cancers

#### Colorectal Cancer (CRC)

CRC is accompanied by an increase in the IL-6 secretion, which also increases with the progression of the cancer, but it is not considered as a prognostic factor of the cancer. IL-6 increases the proliferation and growth of the CRC cells. Proliferation and survival of the cancer cells induced by IL-6 depend on the STAT3 activity in anthrocytes (91, 93). In this type of cancer, IL-6 stimulates the proliferation of the epithelial cells through NF- $\kappa$ B and STAT3 cascading pathways (88).

#### Gastric Cancer (GC)

STAT3 activity is associated with the reduced survival (viability) in the patients with gastric cancer. Studies have shown that the IL-6 activity and function are associated with the enhanced initiating factors of GC. IL-6 is the ligand-binding receptor of gp130 and strongly and effectively upregulates gp130 in the gastric tumors (91).

#### Liver Cancer

HCC, as a common liver cancer is often caused by the inflammation due to the chronic infection with the hepatitis C and B viruses. IL-6 is strongly associated with HCC, and the higher the serum level of IL-6, the larger the size of the tumor. IL-6R and gp130 are highly expressed in the hepatocytes (91).

#### Pancreatic Cancer

Increased STAT3 activity has been proven in patients with pancreatic cancer. Serum level of IL-6 is higher in patients with pancreatic cancer than the normal individuals. STAT3-activated MMP7 is required for tumor progression but is dependent on other STAT3 targets for tumor initiation (91).

#### Prostate Cancer

IL-6 and its receptor are highly expressed in PC. STAT3 activity is inversely related to the progression and metastasis of PC (91).

#### Kidney and Bladder Cancers

IL-6 level is related to the tumor size, stage, and tumor progression, and it is also a factor in metastasis of the kidney and bladder tumors (91). The autocrine effect of IL-6 in kidney cancer is associated with P53 (94).

### 5.4. The Effects of BPA on the TNF- $\alpha$ Secretion from the Adipose Tissue

TNF $\alpha$  is a potent cytokine with an essential role in the intrinsic immunity and also a transmembrane protein with 26KDa molecular weight that is expressed in a wide range of cells, in addition to the macrophages including the adipose tissue as well as the breast, colon, and pancreatic cancer cells. TNF $\alpha$  has two types of receptors (92, 95-97):

- 1) R1 (also called as P55)
- 2) R2 (also called as P75)

R1 is expressed in all cells, and R2 is selectively expressed in the endothelial and immune system cells. TNF $\alpha$  has the same tendency for binding to both receptors. TNF $\alpha$  is essential for the proliferation and proper functioning of the T cells, B cells, macrophages, and dendritic cells (92, 95, 96).

Studies on the effect of BPA on the TNF $\alpha$  secretion from the adipose tissue have proven that BPA increases the expression and secretion of TNF $\alpha$  from the adipose tissue cells through binding to the classic ERs present on the surface of the adipose tissue (30, 98). Lack of TNF $\alpha$  regulation is associated with the extensive changes, such as the development of cancer. TNF $\alpha$  is associated with all stages of cancer, such as tumorigenesis, cellular changes, survival, proliferation, invasion, angiogenesis, and metastasis (95). Most chronic inflammatory diseases increase the risk of cancer, and it has also been proven that most cancers develop in the chronic inflammatory sites. TNF $\alpha$  is not usually detectable in the plasma of healthy individuals, for example, in patients with PC, TNF $\alpha$  and IL-6 levels are associated with the cancer progression (96). Tumor cells that have high growth and proliferation and migrate rapidly induce the TNF $\alpha$  expression, and this cytokine increases the expression of the Vascular Cell Adhesion Molecule-1 (VCAM1). As a result, communication and interaction is established between the tumor and endothelial cells resulting in the colonization in the tissues and tumor metastasis to other tissues (99, 100). TNF $\alpha$  induces the expression of the MMPs and dipeptidyl peptidases and causes the growth, migration, and invasion of ERs in the MCF7 and MDA-MB231 cell lines (101). MMPs are the enzymes that regulate the extracellular matrix and play a role in the angiogenesis, growth, proliferation, and metastasis of various tumors including the BC cells. Epithelial-mesenchymal transmission is among the other roles of TNF $\alpha$  in the progression of cancer. During this process, epithelial cells lose their phenotypic properties and increase the mesenchymal properties such as loss of the E-cadherin (epithelial marker), induction of

vimentin, loss of the cellular adhesion, disruption of the cell adhesion, resistance to apoptosis, and increased mobility and invasion (101).

#### 5.4.1. Increased Level of TNF- $\alpha$ and its Association with Various Cancers

##### Kidney Cancer

TNF $\alpha$  and TNFR2 are expressed in the renal malignancies. Serum level of TNF $\alpha$  is significantly higher in patients with kidney cancer than the healthy individuals and increases with the progression of the cancer (102). TNF $\alpha$  level is associated with the tumor size in kidney cancer. TNF $\alpha$  exacerbates the EMT (epithelial-mesenchymal transitional) expression and tumorigenesis in the RCC through suppression of the E-cadherin, regulation of the vimentin expression and MMP9, and increased invasion to other tissues (103).

##### Breast Cancer (BC)

Breast cancer often develops after a chronic inflammation and in the presence of the immune cells, such as neutrophils, macrophages, TNF $\alpha$ , lymphocyte B, and so on. TNF $\alpha$  is not detectable in the serum of the healthy women, but in the women with BC, the level of cytokines, especially the TNF $\alpha$  is elevated and measurable (103).

##### Pancreatic Cancer

Pancreatic cancer is one of the fatal cancers and the patients with pancreatic cancer do not survive more than 5 years. Pre-inflammatory cytokines as well as the IL-6, IL-17, and TNF $\alpha$  play an important role in the pancreatic cancer. Pancreatic tumor cells can also produce the TNF $\alpha$  at picograms inducing the expression of the EGFR and Transforming Growth Factor alpha (TGF $\alpha$ ), as the mediators for proliferation (93, 96).

## Conclusion

BPA is a synthetic chemical that is increasingly found in the human living environment. This compound has a great effect on the signaling of the adipose tissue through the ERs, which can lead to the changes in the level of the adipocytokines including TNF- $\alpha$ , IL-6, leptin, and APN. Imbalance in the secretion of these adipocytokines can lead the cells to be cancerous. Findings of the existing researches showed that BPA increases the secretion of TNF- $\alpha$ , IL-6, and leptin through various signaling pathways and in contrast, it reduces the level of APN. Given the protective role of the APN, targeting pathways increasing the level of this adipokine will be effective in reducing the side effects of BPA, which requires to be considered in environmental investigation in the further researches.

## Acknowledgments

None.

## Conflict of Interest

The authors confirm that there are no conflicts of interest.

## References

1. Marino M, Pellegrini M, La Rosa P, Acconcia F. Susceptibility of estrogen receptor rapid responses to xenoestrogens: Physiological outcomes. *Steroids*. 2012;77(10):910-7. [DOI:10.1016/j.steroids.2012.02.019] [PMID]
2. Geens T, Aerts D, Berthot C, et al. A review of dietary and non-dietary exposure to bisphenol-A. *Food Chem Toxicol*. 2012;50(10):3725-40. [DOI:10.1016/j.fct.2012.07.059] [PMID]
3. Staples CA, Dorn PB, Klecka GM, O'Block ST, Harris LR. A review of the environmental fate, effects, and exposures of bisphenol A. *Chemosphere*. 1998 Apr;36(10):2149-73. [DOI:10.1016/S0045-6535(97)10133-3]
4. Ma Y, Liu H, Wu J, et al. The adverse health effects of bisphenol A and related toxicity mechanisms. *Environ Res*. 2019;176:108575. [DOI:10.1016/j.envres.2019.108575] [PMID]
5. Welshons WV, Nagel SC, vom Saal FS. Large effects from small exposures. III. Endocrine mechanisms mediating effects of bisphenol A at levels of human exposure. *Endocrinol*. 2006;147(6 Suppl):S56-69. [DOI:10.1210/en.2005-1159] [PMID]
6. Beausoleil C, Emond C, Cravedi JP, et al. Regulatory identification of BPA as an endocrine disruptor: Context and methodology. *Mol Cell Endocrinol*. 2018; 475 (5):4-9. [DOI:10.1016/j.mce.2018.02.001] [PMID]
7. Crain DA, Eriksen M, Iguchi T, et al. An ecological assessment of bisphenol-A: evidence from comparative biology. *Reprod Toxicol*. 2007;24(2):225-39. [DOI:10.1016/j.reprotox.2007.05.008] [PMID]
8. Abraham A, Chakraborty P. A review on sources and health impacts of bisphenol A. *Rev Environ Health*. 2020;35(2):201-10. [DOI:10.1515/reveh-2019-0034] [PMID]
9. Shin BS, Kim CH, Jun YS, et al. Physiologically based pharmacokinetics of bisphenol A. *J Toxicol Environ Health A*. 2004;67(23-24):1971-85. [DOI:10.1080/15287390490514615] [PMID]
10. Hines CJ, Christianson AL, Jackson MV, et al. An evaluation of the relationship among urine, air, and hand measures of exposure to bisphenol A (BPA) in US manufacturing workers. *Ann Work Expo Health*. 2018; 62(7):840-51.

- [DOI:10.1093/annweh/wxy042] [PMID] [PMCID]
11. Wang M, Rang O, Liu F, et al. A systematic review of metabolomics biomarkers for Bisphenol A exposure. *Metabolomics*. 2018;14(4):1-9. [DOI:10.1007/s11306-018-1342-z] [PMID]
  12. Ho KL, Yuen KK, Yau MS, et al. Glucuronide and sulfate conjugates of bisphenol A: Chemical synthesis and correlation between their urinary levels and plasma bisphenol A content in voluntary human donors. *Arch Environ Contam Toxicol*. 2017;73(3):410-20. [DOI:10.1007/s00244-017-0438-1] [PMID]
  13. Thayer KA, Doerge DR, Hunt D, et al. Pharmacokinetics of bisphenol A in humans following a single oral administration. *Environ Int*. 2015;83:107-15. [DOI:10.1016/j.envint.2015.06.008] [PMID] [PMCID]
  14. Pelch KE, Li Y, Perera L, Thayer KA, Korach KS. Characterization of estrogenic and androgenic activities for bisphenol A-like chemicals (BPs): in vitro estrogen and androgen receptors transcriptional activation, gene regulation, and binding profiles. *Toxicol Sci*. 2019;172(1):23-37. [DOI:10.1093/toxsci/kfz173] [PMID] [PMCID]
  15. De Coster S, Van Larebeke N. Endocrine-disrupting chemicals: associated disorders and mechanisms of action. *J Environ Public Health*. 2012;2012. [DOI:10.1155/2012/713696] [PMID] [PMCID]
  16. Pereira-Fernandes A, Demaegdts H, Vandermeiren K, et al. Evaluation of a screening system for obesogenic compounds: screening of endocrine disrupting compounds and evaluation of the PPAR dependency of the effect. *PloS One*. 2013;8(10):e77481. [DOI:10.1371/journal.pone.0077481] [PMID] [PMCID]
  17. Kuiper GG, Gustafsson JÅ. The novel estrogen receptor- $\beta$  subtype: potential role in the cell- and promoter-specific actions of estrogens and anti-estrogens. *FEBS Lett*. 1997;410(1):87-90. [DOI:10.1016/S0014-5793(97)00413-4]
  18. Ben-Jonathan N, Hugo ER, Brandebourg TD. Effects of bisphenol A on adipokine release from human adipose tissue: Implications for the metabolic syndrome. *Mol Cell Endocrinol*. 2009;304(1-2):49-54. [DOI:10.1016/j.mce.2009.02.022] [PMID] [PMCID]
  19. Acconcia F, Pallottini V, Marino M. Molecular mechanisms of action of BPA. DOSE-RESPONSE. 2015;13(4):1559325815610582. [DOI:10.1177/1559325815610582] [PMID] [PMCID]
  20. Wozniak AL, Bulayeva NN, Watson CS. Xenestrogens at picomolar to nanomolar concentrations trigger membrane estrogen receptor- $\alpha$ -mediated Ca<sup>2+</sup> fluxes and prolactin release in GH3/B6 pituitary tumor cells. *Environ Health Perspect*. 2005;113(4):431-9. [DOI:10.1289/ehp.7505] [PMID] [PMCID]
  21. Prossnitz ER, Arterburn JB, Smith HO, Oprea TI, Sklar LA, Hathaway HJ. Estrogen signaling through the transmembrane G protein-coupled receptor GPR30. *Annu Rev Physiol*. 2008;70:165-90. [DOI:10.1146/annurev.physiol.70.113006.100518] [PMID]
  22. Nesan D, Sewell LC, Kurrasch DM. Opening the black box of endocrine disruption of brain development: Lessons from the characterization of Bisphenol A. *Horm Behav*. 2018;101:50-8. [DOI:10.1016/j.yhbeh.2017.12.001] [PMID]
  23. Sohoni P, Sumpter JP. Several environmental oestrogens are also anti-androgens. *J Endocrinol*. 1998;158(3):327-40. [DOI:10.1677/joe.0.1580327] [PMID]
  24. Pellegrini M, Accoccia F, Marino M. Endocrine disruptors. a gender affair. *Open Access Biol*. 2013;1:5.
  25. Ariemma F, D'Esposito V, Liguoro D, et al. Low-dose bisphenol-A impairs adipogenesis and generates dysfunctional 3T3-L1 adipocytes. *PLoS One*. 2016;11(3):e0150762. [DOI:10.1371/journal.pone.0150762] [PMID] [PMCID]
  26. Zoeller RT. New insights into thyroid hormone action in the developing brain: the importance of T3 degradation. *Endocrinol*. 2010;151(11):5089-91. [DOI:10.1210/en.2010-0926] [PMID]
  27. Tabb MM, Blumberg B. New modes of action for endocrine-disrupting chemicals. *Mol Endocrinol*. 2006;20(3):475-82. [DOI:10.1210/me.2004-0513] [PMID]
  28. Sui Y, Ai N, Park SH, et al. Bisphenol A and its analogues activate human pregnane X receptor. *Environ Health Perspect*. 2012;120(3):399-405. [DOI:10.1289/ehp.1104426] [PMID] [PMCID]
  29. Abel J, Haarmann-Stemmann T. An introduction to the molecular basics of aryl hydrocarbon receptor biology. *Biol Chem*. 2010;391(11):1235-48. [DOI:10.1515/bc.2010.128] [PMID]
  30. Rubin BS, Soto AM. Bisphenol A: perinatal exposure and body weight. *Mol Cell Endocrinol*. 2009;304(1-2):55-62.

- [[DOI:10.1016/j.mce.2009.02.023](https://doi.org/10.1016/j.mce.2009.02.023)] [[PMID](#)]  
[[PMCID](#)]
31. Le Corre L, Besnard P, Chagnon MC. BPA, an energy balance disruptor. *Crit Rev Food Sci Nutr.* 2015;55(6):769-77.  
[[DOI:10.1080/10408398.2012.678421](https://doi.org/10.1080/10408398.2012.678421)]  
[[PMID](#)]
  32. Hugo ER, Brandebourg TD, Woo JG, Loftus J, Alexander JW, Ben-Jonathan N. Bisphenol A at environmentally relevant doses inhibits adiponectin release from human adipose tissue explants and adipocytes. *Environ Health Perspect.* 2008;116(12):1642-7.  
[[DOI:10.1289/ehp.11537](https://doi.org/10.1289/ehp.11537)] [[PMID](#)] [[PMCID](#)]
  33. Pallottini V, Bulzomi P, Galluzzo P, Martini C, Marino M. Estrogen regulation of adipose tissue functions: involvement of estrogen receptor isoforms. *Infect Disord Drug Targets.* 2008;8(1):52-60.  
[[DOI:10.2174/187152608784139631](https://doi.org/10.2174/187152608784139631)] [[PMID](#)]
  34. Newell-Fugate AE. The role of sex steroids in white adipose tissue adipocyte function. *Reproduction.* 2017;153(4):R133-49.  
[[DOI:10.1530/REP-16-0417](https://doi.org/10.1530/REP-16-0417)] [[PMID](#)]
  35. Vom Saal FS, Nagel SC, Coe BL, Angle BM, Taylor JA. The estrogenic endocrine disrupting chemical bisphenol A (BPA) and obesity. *Mol Cell Endocrinol.* 2012;354(1-2):74-84.  
[[DOI:10.1016/j.mce.2012.01.001](https://doi.org/10.1016/j.mce.2012.01.001)] [[PMID](#)]  
[[PMCID](#)]
  36. Cooke PS, Heine PA, Taylor JA, Lubahn DB. The role of estrogen and estrogen receptor- $\alpha$  in male adipose tissue. *Mol Cell Endocrinol.* 2001;178(1-2):147-54.  
[[DOI:10.1016/S0303-7207\(01\)00414-2](https://doi.org/10.1016/S0303-7207(01)00414-2)]
  37. Muraki K, Okuya S, Tanizawa Y. Estrogen receptor  $\alpha$  regulates insulin sensitivity through IRS-1 tyrosine phosphorylation in mature 3T3-L1 adipocytes. *Endocrine J.* 2006;0609250044-0609250044. [[DOI:10.1507/endocrj.K06-005](https://doi.org/10.1507/endocrj.K06-005)]  
[[PMID](#)]
  38. Garcia Dos Santos E, Dieudonne MN, Pecquery R, Le Moal V, Giudicelli Y, Lacasa D. Rapid nongenomic E2 effects on p42/p44 MAPK, activator protein-1, and cAMP response element binding protein in rat white adipocytes. *Endocrinol.* 2002;143(3):930-40.  
[[DOI:10.1210/endo.143.3.8678](https://doi.org/10.1210/endo.143.3.8678)] [[PMID](#)]
  39. Dieudonne MN, Leneuve MC, Giudicelli Y, Pecquery R. Evidence for functional estrogen receptors  $\alpha$  and  $\beta$  in human adipose cells: regional specificities and regulation by estrogens. *Am J Physiol Cell Physiol.* 2004;286(3):C655-61.  
[[DOI:10.1152/ajpcell.00321.2003](https://doi.org/10.1152/ajpcell.00321.2003)] [[PMID](#)]
  40. Lee MJ, Lin H, Liu CW, et al. Octylphenol stimulates resistin gene expression in 3T3-L1 adipocytes via the estrogen receptor and extracellular signal-regulated kinase pathways. *Am J Physiol Cell Physiol.* 2008;294(6):C1542-51. [[DOI:10.1152/ajpcell.00403.2007](https://doi.org/10.1152/ajpcell.00403.2007)] [[PMID](#)]
  41. Angle BM, Do RP, Ponzi D, et al. Metabolic disruption in male mice due to fetal exposure to low but not high doses of bisphenol A (BPA): evidence for effects on body weight, food intake, adipocytes, leptin, adiponectin, insulin and glucose regulation. *Reprod Toxicol.* 2013;42:256-68. [[DOI:10.1016/j.reprotox.2013.07.017](https://doi.org/10.1016/j.reprotox.2013.07.017)]  
[[PMID](#)] [[PMCID](#)]
  42. Kim JH, Cho HT, Kim YJ. The role of estrogen in adipose tissue metabolism: insights into glucose homeostasis regulation. *Endocrine J.* 2014:EJ14-0262.
  43. Cust AE, Kaaks R, Friedenreich C, et al. Plasma adiponectin levels and endometrial cancer risk in pre- and postmenopausal women. *J Clin Endocrinol Metab.* 2007;92(1):255-63.  
[[DOI:10.1210/jc.2006-1371](https://doi.org/10.1210/jc.2006-1371)] [[PMID](#)]
  44. Gulen ST, Karadag F, Karul AB, et al. Adipokines and systemic inflammation in weight-losing lung cancer patients. *Lung.* 2012;190(3):327-32.  
[[DOI:10.1007/s00408-011-9364-6](https://doi.org/10.1007/s00408-011-9364-6)] [[PMID](#)]
  45. Jasinski-Bergner S, Kielstein H. Adipokines regulate the expression of tumor-relevant MicroRNAs. *Obes Facts.* 2019;12(2):211-25.  
[[DOI:10.1159/000496625](https://doi.org/10.1159/000496625)] [[PMID](#)] [[PMCID](#)]
  46. Di Zazzo E, Polito R, Bartollino S, et al. Adiponectin as link factor between adipose tissue and cancer. *Int J Mol Sci.* 2019;20(4):839.  
[[DOI:10.3390/ijms20040839](https://doi.org/10.3390/ijms20040839)] [[PMID](#)] [[PMCID](#)]
  47. Avgerinos KI, Spyrou N, Mantzoros CS, Dalamaga M. Obesity and cancer risk: Emerging biological mechanisms and perspectives. *Metabolism.* 2019;92:121-35.  
[[DOI:10.1016/j.metabol.2018.11.001](https://doi.org/10.1016/j.metabol.2018.11.001)] [[PMID](#)]
  48. Miyawaki J, Sakayama K, Kato H, Yamamoto H, Masuno H. Perinatal and postnatal exposure to bisphenol a increases adipose tissue mass and serum cholesterol level in mice. *J Atheroscler Thromb.* 2007;0710100002-0710100002.  
[[DOI:10.5551/jat.E486](https://doi.org/10.5551/jat.E486)] [[PMID](#)]
  49. Hug C, Wang J, Ahmad NS, Bogan JS, Tsao TS, Lodish HF. T-cadherin is a receptor for hexameric and high-molecular-weight forms of Acrp30/adiponectin. *Proc Natl Acad Sci U S A.* 2004;101(28):10308-13.  
[[DOI:10.1073/pnas.0403382101](https://doi.org/10.1073/pnas.0403382101)] [[PMID](#)]  
[[PMCID](#)]
  50. Katira A, Tan PH. Adiponectin and its receptor signaling: an anti-cancer therapeutic target and its

- implications for anti-tumor immunity. *Expert Opin Ther Targets*. 2015;19(8):1105-25. [DOI:10.1517/14728222.2015.1035710] [PMID]
51. Kelesidis I, Kelesidis T, Mantzoros C. Adiponectin and cancer: a systematic review. *Br J Cancer*. 2006;94(9):1221-5. [DOI:10.1038/sj.bjc.6603051] [PMID] [PMCID]
  52. Dieudonne MN, Bussiere M, Dos Santos E, Leneuve MC, Giudicelli Y, Pecquery R. Adiponectin mediates antiproliferative and apoptotic responses in human MCF7 breast cancer cells. *Biochem Biophys Res Commun*. 2006;345(1):271-9. [DOI:10.1016/j.bbrc.2006.04.076] [PMID]
  53. Van Sinderen ML, Steinberg GR, Jrgensen SB, et al. Effects of estrogens on adipokines and glucose homeostasis in female aromatase knockout mice. *PloS One*. 2015;10(8):e0136143. [DOI:10.1371/journal.pone.0136143] [PMID] [PMCID]
  54. Watson CS, Bulayeva NN, Wozniak AL, Alyea RA. Xenoestrogens are potent activators of nongenomic estrogenic responses. *Steroids*. 2007;72(2):124-34. [DOI:10.1016/j.steroids.2006.11.002] [PMID] [PMCID]
  55. Fenichel P, Chevalier N, Brucker-Davis F. Bisphenol A: an endocrine and metabolic disruptor. *Ann Endocrinol (Paris)*. 2013;74(3):211-20. [DOI:10.1016/j.ando.2013.04.002] [PMID]
  56. Ropero AB, Alonso-Magdalena P, Ripoll C, Fuentes E, Nadal A. Rapid endocrine disruption: environmental estrogen actions triggered outside the nucleus. *J Steroid Biochem Mol Biol*. 2006;102(1-5):163-9. [DOI:10.1016/j.jsbmb.2006.09.019] [PMID]
  57. Trujillo ME, Scherer PE. Adiponectin-journey from an adipocyte secretory protein to biomarker of the metabolic syndrome. *J Intern Med*. 2005;257(2):167-75. [DOI:10.1111/j.1365-2796.2004.01426.x] [PMID]
  58. Wright HM, Clish CB, Mikami T, et al. A synthetic antagonist for the peroxisome proliferator-activated receptor gamma inhibits adipocyte differentiation. *J Biol Chem*. 2000;275(3):1873-7. [DOI:10.1074/jbc.275.3.1873] [PMID]
  59. Hiroi T, Okada K, Imaoka S, Osada M, Funae Y. Bisphenol A binds to protein disulfide isomerase and inhibits its enzymatic and hormone-binding activities. *Endocrinol*. 2006;147(6):2773-80. [DOI:10.1210/en.2005-1235] [PMID]
  60. Lin SY, Huang SC, Sheu WH. Circulating adiponectin concentrations were related to free thyroxine levels in thyroid cancer patients after thyroid hormone withdrawal. *Metabolism*. 2010;59(2):195-9. [DOI:10.1016/j.metabol.2009.06.032] [PMID]
  61. Tworoger SS, Eliassen AH, Kelesidis T, et al. Plasma adiponectin concentrations and risk of incident breast cancer. *J Clin Endocrinol Metab*. 2007;92(4):1510-6. [DOI:10.1210/jc.2006-1975] [PMID]
  62. Soliman PT, Wu D, Tortolero Luna G, et al. Association between adiponectin, insulin resistance, and endometrial cancer. *Cancer*. 2006;106(11):2376-81. [DOI:10.1002/cncr.21866] [PMID]
  63. Perrier S, Jarde T. Adiponectin, an anti-carcinogenic hormone? A systematic review on breast, colorectal, liver and prostate cancer. *Curr Med Chem*. 2012;19(32):5501-12. [DOI:10.2174/092986712803833137] [PMID]
  64. Petridou ET, Mitsiades N, Gialamas S, et al. Circulating adiponectin levels and expression of adiponectin receptors in relation to lung cancer: two case-control studies. *Oncology*. 2007;73(3-4):261-9. [DOI:10.1159/000127424] [PMID]
  65. Bub JD, Miyazaki T, Iwamoto Y. Adiponectin as a growth inhibitor in prostate cancer cells. *Biochem Biophys Res Commun*. 2006;340(4):1158-66. [DOI:10.1016/j.bbrc.2005.12.103] [PMID]
  66. Li G, Cong L, Gasser J, Zhao J, Chen K, Li F. Mechanisms underlying the anti-proliferative actions of adiponectin in human breast cancer cells, MCF7-dependency on the cAMP/protein kinase-A pathway. *Nutr Cancer*. 2011;63(1):80-8. [DOI:10.1080/01635581.2010.516472] [PMID]
  67. Yildirim A, Bilici M, Cayir K, Yanmaz V, Yildirim S, Tekin SB. Serum adiponectin levels in patients with esophageal cancer. *Jpn J Clin Oncol*. 2009 ;39(2):92-6. [DOI:10.1093/jjco/hyn143] [PMID]
  68. Hebbard L, Ranscht B. Multifaceted roles of adiponectin in cancer. *Best Pract Res Clin Endocrinol Metab*. 2014;28(1):59-69. [DOI:10.1016/j.beem.2013.11.005] [PMID] [PMCID]
  69. VanSaun MN. Molecular pathways: adiponectin and leptin signaling in cancer. *Clin Cancer Res*. 2013;19(8):1926-32. [DOI:10.1158/1078-0432.CCR-12-0930] [PMID] [PMCID]
  70. Mitsiades N, Pazaitou-Panayiotou K, Aronis KN, et al. Circulating adiponectin is inversely associated with risk of thyroid cancer: in vivo and

- in vitro studies. *J Clin Endocrinol Metab.* 2011;96(12):E2023-8. [[DOI:10.1210/jc.2010-1908](https://doi.org/10.1210/jc.2010-1908)] [[PMID](#)] [[PMCID](#)]
71. Byeon JS, Jeong JY, Kim MJ, et al. Adiponectin and adiponectin receptor in relation to colorectal cancer progression. *Int J Cancer.* 2010;127(12):2758-67. [[DOI:10.1002/ijc.25301](https://doi.org/10.1002/ijc.25301)] [[PMID](#)]
  72. Spyridopoulos TN, Petridou ET, Skalkidou A, Dessypris N, Chrousos GP, Mantzoros CS. Low adiponectin levels are associated with renal cell carcinoma: a case-control study. *Int J Cancer.* 2007;120(7):1573-8. [[DOI:10.1002/ijc.22526](https://doi.org/10.1002/ijc.22526)] [[PMID](#)]
  73. Maffei M, Fei H, Lee GH, et al. Increased expression in adipocytes of ob RNA in mice with lesions of the hypothalamus and with mutations at the db locus. *Proc Natl Acad Sci U S A.* 1995;92(15):6957-60. [[DOI:10.1073/pnas.92.15.6957](https://doi.org/10.1073/pnas.92.15.6957)] [[PMID](#)] [[PMCID](#)]
  74. Vuletic MS, Milosevic VS, Jancic SA, Zujovic JT, Krstic MS, Vukmirovic FC. Clinical significance of Leptin receptor (LEPR) and Endoglin (CD105) expressions in colorectal adenocarcinoma. *J BUON.* 2019;24(6):2448-57.
  75. Volberg V, Harley K, Calafat AM, et al. Maternal bisphenol a exposure during pregnancy and its association with adipokines in Mexican-American children. *Environ Mol Mutagen.* 2013;54(8):621-8.78. [[DOI:10.1002/em.21803](https://doi.org/10.1002/em.21803)] [[PMID](#)] [[PMCID](#)]
  76. Uddin S, Bu R, Ahmed M, et al. Overexpression of leptin receptor predicts an unfavorable outcome in Middle Eastern ovarian cancer. *Mol Cancer.* 2009;8(1):74.79. [[DOI:10.1186/1476-4598-8-74](https://doi.org/10.1186/1476-4598-8-74)] [[PMID](#)] [[PMCID](#)]
  77. Li F, Cao Y, Li J, et al. The clinical significance of serum adipocytokines level in patients with lung cancer. *J Thorac Dis.* 2019;11(8):3547-55.80. [[DOI:10.21037/jtd.2019.07.66](https://doi.org/10.21037/jtd.2019.07.66)] [[PMID](#)] [[PMCID](#)]
  78. Chen C, Chang YC, Liu CL, Chang KJ, Guo C. Leptin-induced growth of human ZR-75-1 breast cancer cells is associated with up-regulation of cyclin D1 and c-Myc and down-regulation of tumor suppressor p53 and p21 WAF1/CIP1. *Breast Cancer Res Treat.* 2006;98(2):121-32.81. [[DOI:10.1007/s10549-005-9139-y](https://doi.org/10.1007/s10549-005-9139-y)] [[PMID](#)]
  79. Barrea L, Gallo M, Ruggeri RM, et al. Nutritional status and follicular-derived thyroid cancer: An update. *Crit Rev Food Sci Nutr.* 2021;61(1):25-59.82. [[DOI:10.1080/10408398.2020.1714542](https://doi.org/10.1080/10408398.2020.1714542)] [[PMID](#)]
  80. Li F, Zhao S, Guo T, Li J, Gu C. The nutritional cytokine leptin promotes NSCLC by activating the PI3K/AKT and MAPK/ERK pathways in NSCLC cells in a paracrine manner. *Biomed Res Int.* 2019;2019:2585743.83. [[DOI:10.1155/2019/2585743](https://doi.org/10.1155/2019/2585743)] [[PMID](#)] [[PMCID](#)]
  81. Machinal-Quélin F, Dieudonné MN, Pecquery R, Leneveu MC, Giudicelli Y. Direct in vitro effects of androgens and estrogens on ob gene expression and leptin secretion in human adipose tissue. *Endocrine.* 2002;18(2):179-84.84. [[DOI:10.1385/ENDO:18:2:179](https://doi.org/10.1385/ENDO:18:2:179)]
  82. Chevalier N, Fénichel P. Bisphenol A: Targeting metabolic tissues. *Rev Endocr Metab Disord.* 2015;16(4):299-309.85. [[DOI:10.1007/s11154-016-9333-8](https://doi.org/10.1007/s11154-016-9333-8)] [[PMID](#)]
  83. Ptak A, Rak-Mardyla A, Gregoraszczyk EL. Cooperation of bisphenol A and leptin in inhibition of caspase-3 expression and activity in OVCAR-3 ovarian cancer cells. *Toxicol In Vitro.* 2013;27(6):1937-43.86. [[DOI:10.1016/j.tiv.2013.06.017](https://doi.org/10.1016/j.tiv.2013.06.017)] [[PMID](#)]
  84. Ptak A, Gregoraszczyk EL. Bisphenol A induces leptin receptor expression, creating more binding sites for leptin, and activates the JAK/Stat, MAPK/ERK and PI3K/Akt signalling pathways in human ovarian cancer cell. *Toxicol Lett.* 2012 ;210(3):332-7.87. [[DOI:10.1016/j.toxlet.2012.02.003](https://doi.org/10.1016/j.toxlet.2012.02.003)] [[PMID](#)]
  85. Hoffmann M, Fiedor E, Ptak A. Bisphenol A and its derivatives tetrabromobisphenol A and tetrachlorobisphenol A induce apelin expression and secretion in ovarian cancer cells through a peroxisome proliferator-activated receptor gamma-dependent mechanism. *Toxicol Lett.* 2017;269:15-22.88. [[DOI:10.1016/j.toxlet.2017.01.006](https://doi.org/10.1016/j.toxlet.2017.01.006)] [[PMID](#)]
  86. Tsai CF, Chen JH, Wu CT, Chang PC, Wang SL, Yeh WL. Induction of osteoclast-like cell formation by leptin-induced soluble intercellular adhesion molecule secreted from cancer cells. *Ther Adv Med Oncol.* 2019;11:1758835919846806.89. [[DOI:10.1177/1758835919846806](https://doi.org/10.1177/1758835919846806)] [[PMID](#)] [[PMCID](#)]
  87. Lipsey CC, Harbuzariu A, Daley-Brown D, Gonzalez-perez RR. Oncogenic role of leptin and Notch interleujin-1 leptin crosstalk outcome in cancer. *World J Methodol.* 2016; 26(6):43-55. - [[DOI:10.5662/wjm.v6.i1.43](https://doi.org/10.5662/wjm.v6.i1.43)] [[PMID](#)] [[PMCID](#)]
  88. Kumari N, Dwarakanath BS, Das A, Bhatt AN. Role of interleukin-6 in cancer progression and therapeutic resistance. *Tumour Biol.* 2016;37(9):11553-72. [[DOI:10.1007/s13277-016-5098-7](https://doi.org/10.1007/s13277-016-5098-7)] [[PMID](#)]

89. Schmidt-Arras D, Rose-John S. IL-6 pathway in the liver: from physiopathology to therapy. *J Hepatol.* 2016;64(6):1403-15. [DOI:10.1016/j.jhep.2016.02.004] [PMID]
90. Wang L, Yi T, Kortylewski M, Pardoll DM, Zeng D, Yu H. IL-17 can promote tumor growth through an IL-6-Stat3 signaling pathway. *J Exp Med.* 2009;206(7):1457-64. [DOI:10.1084/jem.20090207] [PMID] [PMCID]
91. Taniguchi K, Karin M. IL-6 and related cytokines as the critical lynchpins between inflammation and cancer. *Semin Immunol.* 2014;26(1):54-74. [DOI:10.1016/j.smim.2014.01.001] [PMID]
92. Padoan A, Plebani M, Basso D. Inflammation and pancreatic cancer: focus on metabolism, cytokines, and immunity. *Int J Mol Sci.* 2019;20(3):676. [DOI:10.3390/ijms20030676] [PMID] [PMCID]
93. Becker C, Fantini MC, Wirtz S, Nikolaev A, Lehr HA, Galle PR, et al. IL-6 signaling promotes tumor growth in colorectal cancer. *Cell Cycle.* 2005;4(2):217-20. [DOI:10.4161/cc.4.2.1413] [PMID]
94. Aggarwal BB, Shishodia S, Sandur SK, Pandey MK, Sethi G. Inflammation and cancer: how hot is the link?. *Biochem Pharmacol.* 2006;72(11):1605-21. [DOI:10.1016/j.bcp.2006.06.029] [PMID]
95. Sethi G, Sung B, Aggarwal BB. TNF: a master switch for inflammation to cancer. *Front biosci.* 2008;13(2):5094-107. [DOI:10.2741/3066] [PMID]
96. Balkwill F. TNF- $\alpha$  in promotion and progression of cancer. *Cancer Metastasis Re.* 2006;25(3):409-16. [DOI:10.1007/s10555-006-9005-3] [PMID]
97. Xu LL, Shi CM, Xu GF, et al. TNF- $\alpha$ , IL-6, and leptin increase the expression of miR-378, an adipogenesis-related microRNA in human adipocytes. *Cell Biochem Biophys.* 2014;70(2):771-6. [DOI:10.1007/s12013-014-9980-x] [PMID]
98. Ahmed RG. Maternal bisphenol A alters fetal endocrine system: Thyroid adipokine dysfunction. *Food Chem Toxicol.* 2016;95:168-74. [DOI:10.1016/j.fct.2016.06.017] [PMID]
99. Kitakata H, Nemoto-Sasaki Y, Takahashi Y, Kondo T, Mai M, Mukaida N. Essential roles of tumor necrosis factor receptor p55 in liver metastasis of intrasplenic administration of colon 26 cells. *Cancer Res.* 2002;62(22):6682-7.
100. Zagozda M, Sarnecka AK, Staszczak Z, Nowak-Niezgoda M, Durlak M. Correlation of TNF- $\alpha$  and TGF- $\beta$  polymorphisms with protein levels in pancreatic ductal adenocarcinoma and colorectal cancer. *Contemp Oncol (Pozn).* 2019;23(4):214-219. [DOI:10.5114/wo.2019.91537] [PMID] [PMCID]
101. Martínez-Reza I, Díaz L, García-Becerra R. Preclinical and clinical aspects of TNF- $\alpha$  and its receptors TNFR1 and TNFR2 in breast cancer. *J Biomed Sci.* 2017;24(1):90. [DOI:10.1186/s12929-017-0398-9] [PMID] [PMCID]
102. Al-Lamki RS, Mayadas TN. TNF receptors: signaling pathways and contribution to renal dysfunction. *Kidney Int.* 2015;87(2):281-96. [DOI:10.1038/ki.2014.285] [PMID]
103. Ho MY, Tang SJ, Chuang MJ, et al. TNF- $\alpha$  induces epithelial-mesenchymal transition of renal cell carcinoma cells via a GSK3 $\beta$ -dependent mechanism. *Mol Cancer Res.* 2012;10(8):1109-19. [DOI:10.1158/1541-7786.MCR-12-0160] [PMID]

#### How to Cite This Article:

Karbalaee Hasani A, Nejad Ebrahimi Z, Mahami N, Alizadeh M, Rasooli Z, Hemmati M. A Review of Bisphenol A Effects on White Adipose Tissue and Its Correlation with Various Cancers. *J Adv Med Biomed Res.* 2022; 30 (139): 86-100.

#### Download citation:

[BibTeX](#) | [RIS](#) | [EndNote](#) | [Medlars](#) | [ProCite](#) | [Reference Manager](#) | [RefWorks](#)

#### Send citation to:

 [Mendeley](#)  [Zotero](#)  [RefWorks](#) [RefWorks](#)