

Rheum turkestanicum Janisch Root Extract Mitigates 6-OHDA-Induced Neuronal Toxicity Against Human Neuroblastoma SH-SY5Y Cells

Roghayeh Rashidi¹ , Amir Reza Afshari² , Hamid Mollazadeh³ , Azar Hosseini^{3*} 

1. Dept. of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran
2. Dept. of Physiology and Pharmacology, Faculty of Medicine, North Khorasan University of Medical Sciences, Bojnourd, Iran
3. Pharmacological Research Center of Medicinal Plants, Mashhad University of Medical Sciences, Mashhad, Iran

Article Info

 [10.30699/jambs.30.143.553](https://doi.org/10.30699/jambs.30.143.553)

Received: 2021/05/11;
Accepted: 2022/07/13;
Published Online: 10 Oct 2022;

Use your device to scan and read the article online



Corresponding Information:

Azar Hosseini,

Pharmacological Research Center of Medicinal Plants, Mashhad University of Medical Sciences, Mashhad, Iran

E-Mail: HosseiniAZ@mums.ac.ir

ABSTRACT

Background & Objective: *Rheum turkestanicum* (*R. turkestanicum*) has been known to reduce inflammation and has antioxidant properties such as protective effect in neurons. This study aimed to determine the effects of *R. turkestanicum* on neuronal toxicity induced by the pro-parkinsonian neurotoxin 6-hydroxydopamine (6-OHDA) in neuroblastoma SH-SY5Y cells.

Materials & Methods: MTT and DNA fragmentation by PI staining (sub-G1 peak) assays were used to determine cell viability and induction of apoptosis, respectively. Fluorimetry methods measured lipid peroxidation malondialdehyde (MDA) and reactive oxygen species (ROS) levels. The amount of glutathione (GSH) and the activity of superoxide dismutase (SOD) were measured by DTNB (5, 5'-Dithiobis (2-nitrobenzoic acid) and pyrogallol respectively.

Results: Pretreatment with 12.5 to 100 µg/mL of *R. turkestanicum* extract for 24 hours attenuated 6-OHDA (final concentration 42.5 µg/mL)-induced cytotoxicity. Also, the pretreatment of SH-SY5Y cells with *R. turkestanicum* inhibited 6-OHDA-stimulated apoptosis in a dose-dependent manner. Additionally, *R. turkestanicum* extract repressed 6-OHDA-induced oxidative stress as measured by the MDA, ROS, GSH, and SOD levels.

Conclusion: The findings suggest that *R. turkestanicum* extract has neuroprotective activity on 6-OHDA-induced neuronal toxicity of neuroblastoma cells.

Keywords: *Rheum turkestanicum*, Oxidative stress, 6-Hydroxydopamine, Neuroblastoma, Apoptosis



Copyright © 2022. This is an original open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permits copy and redistribution of the material just in noncommercial usages with proper citation.

Introduction

As the most well-known progressive neurodegenerative disorder, Parkinson's disease (PD), is identified by a selected loss of dopaminergic neurons within the brain (1), which results in debilitating symptoms including uncontrollable tremor, bradykinesia, and inflexibility (2). Supplementation with levodopa has shown efficient therapeutic benefits for the treatment of PD. The reason for idiopathic PD is unclear. Prior surveys have shown that various triggers are involved in the pathogenesis of PD, such as generation of free radicals and dysfunction of mitochondrial and natural agents, which lead to toxicity (3).

The amount of oxidative stress is accumulated in the substantia nigra of PD patients (4). The production of MDA following lipid peroxidation and reactive oxygen species [ROS], environmental toxins, and mitochondrial dysfunction are the leading causes of PD (5). Based on these findings, excessive ROS production could be critically related to the pathology of neurodegenerative

problems (6, 7). 6-hydroxydopamine (6-OHDA) is a toxin that injures neurons and is classically used in neurodegenerative models as it leads to oxidative damage and cell death of dopaminergic cells. Several studies have indicated that modulation of oxidative stress can decrease the degeneration of neurons following 6-OHDA toxicity, which is utilized to develop neurodegenerative laboratory models (8).

Rheum species is related to the Polygonaceae plant family and is used widely in traditional Chinese medicine. Anthraquinones, phenolic compounds, dianthrones, glycosides, and tannins are active compounds of the *Rheum* species. *R. turkestanicum* Janischew grows widely in central Asia and the northeast of Iran (9). The root of *R. turkestanicum* is a well-known traditional Iranian herbal medicine commonly used to treat hypertension, diabetes, and cancer (10). Different investigations have reported that the species of *Rheum*, including *R. undulatum* contains antioxidant compounds

(11-13). It is reported that *R. undulatum* diminishes ROS and hydrogen peroxide (14).

Further studies have indicated that *R. emodi* reduces the toxicity of cardiac cells against H_2O_2 (10, 15). Despite the antioxidant nature of *R. turkestanicum*, no experimental evidence exists to prove the neuroprotective effect of this medicinal herb. Hence, this research evaluated the protective impacts of *R. turkestanicum* hydroalcoholic extract on 6-OHDA-stimulated neuronal damage against SH-SY5Y. The current outcomes revealed that *R. turkestanicum* suppressed 6-OHDA-induced neuronal toxicity by inhibiting free radicals and MDA levels in human neuroblastoma *in vitro*.

Materials and Methods

Reagents

After receiving approval from the Ethics Committee, the experiments on cells were done according to national guidelines (IR.MUMS.MEDICAL.REC.1397.587). MTT powder (98%) was prepared from Thermo Fisher (Thermo Fisher Scientific, Inc., Waltham, MA, USA). 6-hydroxydopamine (6-OHDA), propidium-iodide (PI), the fluorescent probe 2, 7-dichlorofluorescein diacetate (DCF-DA), dimethyl sulfoxide (DMSO), and phosphate buffer saline (PBS) were acquired from Sigma-Aldrich (St. Louis, MO, USA). RNase A was bought from Alfa Aesar (Ward Hill, MA, USA). High glucose Dulbecco Modified Eagle's medium (DMEM) and fetal bovine serum (FBS) were obtained from Gibco (Grand Island, NY). Trichloroacetic acid (TCA), TBA (thiobarbituric acid), and DTNB (5,5'-Dithiobis(2-nitrobenzoic acid)) were prepared from Merck (Darmstadt, Germany).

The preparation of hydroalcoholic extract

The roots of *R. turkestanicum* were acquired from Mashhad, Khorasan Razavi Province, Iran, and it was verified by Joharchi at Ferdowsi University of Mashhad (herbarium number: 21377). Briefly, the dried powder of the roots (50 g) were separated with 70% ethanol in a Soxhlet apparatus for 48 hours. The obtained hydroalcoholic extract was dried in a water bath and then kept at -18°C in a freezer. The yield of the extract was 19% (w/w).

Cell culture and treatments

SH-SY5Y was bought from the National Cell Bank of Iran (NCBI, Tehran, Iran). SH-SY5Y cells were cultured in an incubator in a humidified atmosphere containing CO_2 (5%) and then passaged at 80% confluence. Pretreatments were done by different concentrations of *R. turkestanicum* extract (12.5 to 100 $\mu\text{g/mL}$), and then 6-OHDA was added (0 to 85 $\mu\text{g/mL}$).

Measurement of cytotoxicity

SH-SY5Y cells were cultured at a density of 10^4 cells/well onto flat-bottomed 96-well culture plates. Then, the cells were exposed to a toxic agent (0 to 85 $\mu\text{g/mL}$) for 24 hours. The MTT assay evaluated cytotoxicity, and the final concentration of 6-OHDA was calculated based on the IC_{50} . Subsequently, the cells were treated with 12.5

to 100 $\mu\text{g/mL}$ of the extract for 2 hours and then exposed to 6-OHDA (final concentration 42.5 $\mu\text{g/mL}$) for 24 hours to determine the inhibitory effect of *R. turkestanicum* on 6-OHDA-induced cytotoxicity. Then, MTT solution (0.5 mg/mL) was added to the cells for 4 hours. The absorbance was read using a microplate reader (Start Fax-2100, UK) at 545 nm (630 nm as a reference) (16). All the treatments were done in triplicates.

Measurement of ROS amount

The ROS level was measured by using the 2,7-dichlorofluorescein diacetate (DCFH-DA) method (7). Briefly, after exposure of the cells to various concentrations of the extract (12.5 to 100 $\mu\text{g/mL}$) for 24 hours, 6-OHDA was added to the cells at the IC_{50} dose of (42.5 $\mu\text{g/mL}$). After 24 hours of incubation, the upper-medium was separated, and the cells were incubated with DCFH-DA (10 μM) for 30 min at 37°C in the dark. Next, the cells were washed with PBS, and the fluorescence intensity of DCF was determined using the FACScan fluorescence plate reader (Becton Dickinson, San Jose, USA) with an excitation/emission of 485/535 nm. All the treatments were done in triplicates.

Lipid peroxidation assay

The MDA level was measured as the end product of lipid peroxidation (17). SH-SY5Y cells were pretreated with 25, 50, and 100 $\mu\text{g/mL}$ of the extract for 24 hours before exposure to 42.5 $\mu\text{g/mL}$ 6-OHDA, followed by an additional 24 hours of incubation. Next, the centrifugation of the cells was done at 13,000 g at 4°C for 30 min. Then, 400 μL of TCA (15%) and 800 μL of TBA (0.7%) were added to 500 μL of the cell samples. The mixture was vortexed and then heated for 40 min, and then 200 μL of the sample was transferred to a 96-well plate, and the fluorescence intensity was read with excitation/emission of 480/530 nm. All the treatments were performed in triplicates.

Flow cytometry analysis of apoptosis

The sub-G1 peak of the treated cells was detected by PI staining and flow cytometry to evaluate the number of apoptotic cells (18). SH-SY5Y cells were pretreated with 25, 50, and 100 $\mu\text{g/mL}$ of the extract for 24 hours before exposure to 42.5 $\mu\text{g/mL}$ 6-OHDA, followed by an additional 24 hours of incubation. Next, the cells were incubated at 37°C in the dark with 400 μL of hypotonic buffer (50 $\mu\text{g/mL}$ PI in 0.1% sodium citrate and 0.1% Triton X-100) for 30 min. Then, cell cycle distribution was analyzed from 10^4 cells using a BD FacsCalibur™ flow cytometer (Becton Dickinson, Mountain View, CA, USA). Afterward, the cell cycle analysis of flow cytometry data was done utilizing the software FlowJo® vX.0.7 (Tree Star, Ashland, OR, USA) (19, 20). All the treatments were done in triplicates.

Measurement of GSH level

The content of the sulfhydryl group was evaluated via the formation of yellow color in the presence of DTNB [5, 5'-Dithiobis (2-nitrobenzoic acid)]. In this regard, phosphate-buffered saline was applied to wash the cells, and then 15% TCA was added to extract GSH from the

cells. After centrifugation at 14,000 rpm for 10 min, the denatured proteins were isolated. In brief, TCA extract (500 mL) was mixed with 1 mL of a reaction mixture containing 0.1 M sodium phosphate buffer (pH 7.5) and 0.6 mM DTNB. A spectrophotometer was used to measure the absorbance at 412 nm for 2 min.

Measurement of SOD activity

The effective doses of the extract were added to the cells and then incubated with 250 μ M 6-OHDA for 24 hours to determine SOD activity. The activity was measured by spectrophotometry (at 405 nm) based on the prevention of pyrogallol autoxidation as described previously. The autoxidation of pyrogallol in tris-cacodylic acid buffer (0.05 M, pH 8.2) was measured.

Statistical analysis

All data were displayed as the mean \pm SEM of at least three independent tests. The data were exposed to statistical analysis via one-way ANOVA followed by Dunnett's test with GraphPad Prism[®] 7.01 software

(GraphPad Software, Inc., San Diego, CA). Mean values were considered to be statistically significant at $p < 0.05$.

Results

R. turkestanicum extract attenuates 6-OHDA-induced cytotoxicity

As demonstrated in [Fig. 1A](#), the treatment of SH-SY5Y cells with 6-OHDA for 24 hours remarkably reduced the cell viability concentration dependently ($p < 0.05$ and $p < 0.001$). Moreover, the IC_{50} for 6-OHDA was about 42.5 μ g/mL. Therefore, the treatment of 42.5 μ g/mL 6-OHDA for 24 hours induced SH-SY5Y cellular toxicity in sequential experiments. To evaluate the cytotoxicity of *R. turkestanicum*, we subjected SH-SY5Y cells to 6.25–200 μ g/mL concentrations of *R. turkestanicum* hydroalcoholic extract for 24 hours, as shown in [Figure. 1B](#). The results showed that *R. turkestanicum* extract in concentrations of 6.25 to 200 μ g/mL did not induce any apparent cytotoxicity ($p > 0.05$, [Figure. 1B](#)).




Figure 1. 6-OHDA (A) and *R. turkestanicum* (B) cytotoxicity in SH-SY5Y cells. The MTT assay evaluated *R. turkestanicum* and 6-OHDA cytotoxicity. The IC_{50} of 6-OHDA was determined at about 42.5 μ g/mL for 24 hours of treatment (A). *R. turkestanicum* had no significant change in the survival of SH-SY5Y cells after 48 hours (B). Each column represents the mean \pm SEM for each group. * $P < 0.05$ and *** $P < 0.001$ versus the control group ($n = 8$).

Based on these data, the cells were pretreated with 12.5, 25, 50, and 100 μ g/mL of the extract for 24 hours before exposure to 42.5 μ g/mL 6-OHDA, followed by an additional 24 hours of incubation to test whether *R. turkestanicum* treatment increased cell survival decreased by 6-OHDA. As a result, the treatment of SH-SY5Y cells with 6-OHDA markedly reduced cell viability compared to the control group ($p < 0.001$) for 24 hours. Interestingly, the pretreatment with *R. turkestanicum* extract (12.5, 25, 50, and 100 μ g/mL) significantly attenuated the cytotoxic effects induced by 6-OHDA compared to the 6-OHDA-treated cells ([Figure 2](#), $p < 0.001$).




Figure 2. *R. turkestanicum* extract attenuated 6-OHDA-induced cytotoxicity in SH-SY5Y cells. SH-SY5Y cells were incubated with *R. turkestanicum* (12.5, 25, 50, and 100 μ g/mL) for 24 hours and stimulated with 6-OHDA (42.5 μ g/mL) for 24 hours. Cytotoxicity was determined by utilizing the MTT assay. The tests were repeated three times independently, with one representative result shown. The data are expressed as the mean \pm SEM of five determinations in a representative experiment. *** $p < 0.001$ versus the control group; ### $p < 0.001$ versus the 6-OHDA-treated cells.

R. turkestanicum extract diminishes ROS and MDA production induced by 6-OHDA

To determine whether *R. turkestanicum* extract could suppress 6-OHDA-initiated oxidative stress, we measured the ROS and MDA levels. As shown in [Fig. 3](#), after treatment of the cells with 42.5 µg/mL 6-OHDA, the ROS and MDA levels were substantially expanded compared to the control group ($p < 0.001$), suggesting that 6-OHDA could incite oxidative stress in SH-SY5Y cells.

Interestingly, pretreatment with *R. turkestanicum* extract (6.25, 12.5, 25, and 50 µg/mL) for 24 hours, followed by exposure to 42.5 µg/mL 6-OHDA for 24

hours, decreased the ROS level significantly compared to the 6-OHDA-treated cells ([Fig. 3A](#), $p < 0.05$ and $p < 0.01$).

Also, twenty-four hours after incubation, the level of lipid peroxidation was examined by measuring the MDA level. In [Figure. 3B](#), the exposure of SH-SY5Y cells to 6-OHDA resulted in a significant increase in the MDA level (192.72%, $p < 0.001$) compared to control cells cultured in the absence of 6-OHDA (100%). The MDA content was significantly diminished in cells pretreated with 25 µg/mL (121%, $p < 0.001$), 50 µg/mL (130%, $p < 0.001$), and 100 µg/mL (121%, $p < 0.001$) of *R. turkestanicum* extract compared to the 6-OHDA-treated cells.




Figure 3. (A) *R. turkestanicum* extract suppressed the 6-OHDA-induced ROS level, which induced apoptosis in SH-SY5Y cells. SH-SY5Y cells were incubated with *R. turkestanicum* (6.25, 12.5, 25, and 50 µg/mL) for 24 hours and stimulated with H₂O₂ (200 µM) for 2 hours and 6-OHDA (42.5 µg/mL) for 24 hours. The ROS level was determined to utilize a fluorescence probe, DCFH-DA. The tests were repeated three times independently, with one representative result demonstrated. The data are expressed as the mean ± SEM of five determinations in a representative experiment. *** $p < 0.001$ versus the control group; ## $p < 0.01$ and # $p < 0.05$ versus the 6-OHDA-treated cells. (B) *R. turkestanicum* extract suppressed 6-OHDA-induced MDA production, which induced oxidative damage in SH-SY5Y cells. SH-SY5Y cells were incubated with *R. turkestanicum* (25, 50, and 100 µg/mL) for 24 hours and stimulated with 6-OHDA (42.5 µg/mL) for 24 hours. The tests were repeated three times independently, with one representative result shown. The data are expressed as the mean ± SEM of five determinations in a representative experiment. *** $p < 0.001$ versus the control group; ### $p < 0.001$ versus the 6-OHDA-treated cells.

R. turkestanicum extract suppresses 6-OHDA-induced apoptosis

As shown in [Figure 4](#), following 24 hours of incubation with 42.5 µg/mL 6-OHDA, the outcomes indicated that in 6-OHDA treatment alone, the percentage of apoptotic cells (67.0 %, $p < 0.001$) was expanded markedly compared with the normal control (11.6 %). However, it respectively dropped to 53.1, 48.6, and 28.1% after 25, 50, and 100 µg/mL of *R. turkestanicum* extract treatments.

R. turkestanicum extract increases GSH levels and SOD activity

We measured SOD activity and the GSH level to determine whether *R. turkestanicum* extract could mediate the 6-OHDA-reduced GSH level and SOD activity. As shown in [Fig. 5](#), after treatment of the cells

with 42.5 µg/mL 6-OHDA, the GSH and SOD levels substantially decreased compared to the control group ($p < 0.01$ and $p < 0.001$, respectively). Interestingly, pretreatment with *R. turkestanicum* extract (25, 50, and 100 µg/mL) for 24 hours, followed by exposure to 42.5 µg/mL 6-OHDA for 24 hours, increased the GSH level significantly compared to the 6-OHDA-treated cells ([Fig. 5A](#), $p < 0.001$).

Also, SOD activity was measured 24 hours after incubation. In [Fig. 5B](#), the exposure of SH-SY5Y cells to 6-OHDA resulted in a significant decrease in SOD activity ($p < 0.001$) compared to control cells cultured in the absence of 6-OHDA (100%). The content of SOD activity significantly increased in cells pretreated with 25, 50, and 100 µg/mL of *R. turkestanicum* extract compared to the 6-OHDA-treated cells.




Figure 4. The effect of *R. turkestanicum* extract on 6-OHDA-induced apoptosis in SH-SY5Y cells. SH-SY5Y cells were incubated with different concentrations of *R. turkestanicum* (25, 50, and 100 µg/mL) and 6-OHDA (42.5 µg/mL) for 24 hours. Then, the apoptosis rate was measured with a PI staining. Values are shown as the mean \pm SEM. The tests were repeated three times independently, with one representative result shown. *** $p < 0.001$ versus the control; #### $p < 0.001$ versus the 6-OHDA-treated cells.




Figure 5. (A) *R. turkestanicum* extract increased the 6-OHDA-reduced GSH level. SH-SY5Y cells were incubated with *R. turkestanicum* (25, 50, and 100 µg/mL) for 24 hours and stimulated with 6-OHDA (42.5 µg/mL) for 24 hours. Data are expressed as the mean \pm SEM of five determinations in a representative experiment. ** $p < 0.01$ versus the control group; #### $p < 0.001$ versus the 6-OHDA-treated cells. (B) *R. turkestanicum* extract increased 6-OHDA-reduced SOD activity. SH-SY5Y cells were incubated with *R. turkestanicum* (25, 50, and 100 µg/mL) for 24 hours and stimulated with 6-OHDA (42.5 µg/mL) for 24 hours. The tests were repeated three times independently, with one representative result shown. The data are expressed as the mean \pm SEM of five determinations in a representative experiment. *** $p < 0.001$ versus the control group; #### $p < 0.001$ versus the 6-OHDA-treated cells.

Discussion

The neuropathological and biochemical features of PD can be presented by 6-OHDA, a neurotoxic agent frequently used to explore molecular pathway patterns involved in cytotoxicity, apoptosis, and oxidative stress in human neuroblastoma SH-SY5Y cells. 6-OHDA is oxidized under physiological circumstances, generating a variety of ROS, including p-quinones and hydrogen peroxides, causing damage to the DNA, lipid peroxidation, and cell death (21-23). The present study showed that *R. turkestanicum* effectively inhibited 6-OHDA-induced neuronal toxicity in SH-SY5Y cells.

While numerous studies have reported that medicinal herbs can preserve cells against oxidative stress-induced toxicity (6, 7, 24-28), it remains unclear whether *R. turkestanicum*, one of the promising medicinal plants, can protect neuronal cells against oxidative stress-induced toxicity. *R. turkestanicum*, as a natural medicine utilized in traditional medicine, has polyphenol ingredients like anthraquinones, stilbenes, anthocyanins, and flavonols (12). The majority of its components have antioxidant activities against toxicological conditions. Studies have indicated that this potential plant is utilized to treat diabetes, hypertension, and cancer (10, 13). Hosseini *et al.* demonstrated the protective effects of *R. turkestanicum* against doxorubicin in cardiomyocytes cells (10), although its protective effects on neuronal toxicity have not been elucidated.

Oxidative stress, as a fundamental reason for cell damage in various debilitating diseases, including neurodegenerative disorders, refers to the imbalance between the formation and clearance of free radical species within the cells. Evidence suggests that 6-OHDA-induced ROS generation plays a primary role in the induction of neuronal damage and apoptosis (29). Hence, evacuation of excess ROS or concealment of the ROS level might be valuable in preventing oxidative neuronal cell death (30, 31). The defensive impacts of *R. turkestanicum* on the cytotoxic activity induced by 6-OHDA were examined to determine the mechanism by pretreatment with *R. turkestanicum* against 6-OHDA-induced injury. It was found that cytoprotective action was of particular interest in oxidative stress (32). In the present study, we found that *R. turkestanicum* significantly mitigated 6-OHDA-induced ROS generation and cytotoxicity in SH-SY5Y cells.

Moreover, MDA reflects the oxidative injury of the plasma membrane, and the resulting TBA reactive substances are related to oxidant stress and lipid peroxidation (33). Hosseini and his colleagues showed that *R. turkestanicum* (12-100 µg/mL) reduced H₂O₂-induced oxidative stress by reducing MDA levels (34). We showed that MDA levels decreased after treatment with *R. turkestanicum*. Also, *R. turkestanicum* extract significantly reduced the apoptosis rate due to 6-OHDA concentration-dependently.

Different studies have reported that 6-OHDA utilizes its toxic effects by producing superoxide anions and hydrogen peroxide. The SOD enzyme converts the superoxide radicals to hydrogen peroxide (35-37). Our results showed that 6-OHDA significantly reduced the GSH level and SOD activity while *R. turkestanicum* extract increased these parameters.

The study's results prove that cell oxidative injury plays an essential role in the toxicity of 6-OHDA and that pretreatment with *R. turkestanicum* can attenuate these changes. This study indicated that *R. turkestanicum* significantly diminished 6-OHDA-induced cytotoxicity, apoptosis, and oxidative stress compared to the 6-OHDA treated-cells. According to the presence of tannins and phenolic compounds (gallic acid) in *R. turkestanicum* (13, 38), the protective effect of *R. turkestanicum* is probably related to its components and other ingredients with antioxidant activities.

Conclusion

The current study offers the first evidence of the protective impact of *R. turkestanicum* on neuronal toxicity. The present outcomes showed that *R. turkestanicum*, a potential medicinal herb, markedly suppressed 6-OHDA-induced neuronal toxicity in human neuroblastoma SH-SY5Y cells, presumably through attenuating ROS, generating MDA, apoptosis, and increasing SOD activity and the GSH level. Albeit, further broader examinations of *in vivo* models of neurodegenerative diseases are essential to recognize.

Acknowledgments

None.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- Dickson DW. Neuropathology of Parkinson disease. *Parkinsonism Relat D.* 2018;46:30-3. [DOI:10.1016/j.parkreldis.2017.07.033] [PMID] [PMCID]
- Deng H, Wang P, Jankovic J. The genetics of Parkinson disease. *Ageing Res Rev.* 2018;42:72-85. [DOI:10.1016/j.arr.2017.12.007] [PMID]
- Bose A, Beal MF. Mitochondrial dysfunction and oxidative stress in induced pluripotent stem cell models of Parkinson's disease. *Eur J Neurosci.* 2019;49(4):525-32. [DOI:10.1111/ejn.14264] [PMID]

4. Ouyang M, Shen X. Critical role of ASK1 in the 6-hydroxydopamine-induced apoptosis in human neuroblastoma SH-SY5Y cells. *J Neurochem.* 2006;97(1):234-44. [[DOI:10.1111/j.1471-4159.2006.03730.x](https://doi.org/10.1111/j.1471-4159.2006.03730.x)] [[PMID](#)]
5. Hao C, Qu A, Xu L, et al. Phenylalanine-mediated Porous CuxO nanoparticle clusters ameliorate parkinson's disease by reducing oxidative stress. *J Am Chem Soc.* 2018;141(2):1091-99. [[DOI:10.1021/jacs.8b11856](https://doi.org/10.1021/jacs.8b11856)] [[PMID](#)]
6. Afshari AR, Sadeghnia HR, Mollazadeh H. A review on potential mechanisms of Terminalia chebula in alzheimer's disease. *Adv Pharmacol Pharm Sci.* 2016;2016:8964849. [[DOI:10.1155/2016/8964849](https://doi.org/10.1155/2016/8964849)] [[PMID](#)] [[PMCID](#)]
7. Sadeghnia HR, Jamshidi R, Afshari AR, Mollazadeh H, Forouzanfar F, Rakhshandeh H. Terminalia chebula attenuates quinolinate-induced oxidative PC12 and OLN-93 cell death. *Mult Scler Relat Disord.* 2017;14:60-7. [[DOI:10.1016/j.msard.2017.03.012](https://doi.org/10.1016/j.msard.2017.03.012)] [[PMID](#)]
8. Blum D, Torch S, Nissou MF, Benabid AL, Verna JM. Extracellular toxicity of 6-hydroxydopamine on PC12 cells. *Neurosci Lett.* 2000;283(3):193-6. [[DOI:10.1016/S0304-3940\(00\)00948-4](https://doi.org/10.1016/S0304-3940(00)00948-4)]
9. Mousa-Al-Reza Hadjzadeh ZR, Khodaei E, Malek M, Ghanbari H. Rheum turkestanicum rhizomes possess anti-hypertriglyceridemic, but not hypoglycemic or hepatoprotective effect in experimental diabetes. *Avicenna J Phytomed.* 2017;7(1):1.
10. Hosseini A, Rajabian A. Protective effect of Rheum turkestanicum root against doxorubicin-induced toxicity in H9C2 cells. *Braz J Psychiatry.* 2016;26(3):347-51. [[DOI:10.1016/j.bjp.2016.02.004](https://doi.org/10.1016/j.bjp.2016.02.004)]
11. Hosseini A, Mollazadeh H, Amiri MS, Sadeghnia HR, Ghorbani A. Effects of a standardized extract of Rheum turkestanicum Janischew root on diabetic changes in the kidney, liver and heart of streptozotocin-induced diabetic rats. *Biomed Pharmacother.* 2017;86:605-11. [[DOI:10.1016/j.biopha.2016.12.059](https://doi.org/10.1016/j.biopha.2016.12.059)] [[PMID](#)]
12. Shiezadeh F, Mousavi SH, Amiri MS, Iranshahi M, Tayarani-Najaran Z, Karimi G. Cytotoxic and apoptotic potential of rheum turkestanicum Janischew root extract on human cancer and normal cells. *Iran J Pharm Res.* 2013;12(4):811.
13. Yazdi MET, Khara J, Sadeghnia HR, Bahabadi SE, Darroudi M. Biosynthesis, characterization, and antibacterial activity of silver nanoparticles using Rheum turkestanicum shoots extract. *Res Chem Intermed Res Chem.* 2018;44(2):1325-34. [[DOI:10.1007/s11164-017-3169-z](https://doi.org/10.1007/s11164-017-3169-z)]
14. Zhang R, Kang KA, Piao MJ, et al. Rhapontigenin from rheum undulatum protects against oxidative-stress-induced cell damage through antioxidant activity. *J Toxicol Environ Health Part A.* 2007;70(13):1155-66. [[DOI:10.1080/15287390701252766](https://doi.org/10.1080/15287390701252766)] [[PMID](#)]
15. Chai Y, Wang F, Li Y, Liu K, Xu H. Antioxidant activities of stilbenoids from rheum emodi wall. *Evid Based Complement Alternat Med.* 2012; 2012:603678. [[DOI:10.1155/2012/603678](https://doi.org/10.1155/2012/603678)] [[PMID](#)] [[PMCID](#)]
16. Mousavi SH, Tavakkol-Afshari J, Brook A, Jafari-Anarkooli I. Role of caspases and bax protein in saffron-induced apoptosis in MCF-7 cells. *Food Chem Toxicol.* 2009;47(8):1909-13. [[DOI:10.1016/j.fct.2009.05.017](https://doi.org/10.1016/j.fct.2009.05.017)] [[PMID](#)]
17. Mollazadeh H, Boroushaki MT, Soukhtanloo M, Afshari AR, Vahedi MM. Effects of pomegranate seed oil on oxidant/antioxidant balance in heart and kidney homogenates and mitochondria of diabetic rats and high glucose-treated H9c2 cell line. *Avicenna J Phytomed.* 2017;7(4):317.
18. Riccardi C, Nicoletti I. Analysis of apoptosis by propidium iodide staining and flow cytometry. *Nat Protoc.* 2006;1(3):1458. [[DOI:10.1038/nprot.2006.238](https://doi.org/10.1038/nprot.2006.238)] [[PMID](#)]
19. Tayarani-Najaran Z, Sareban M, Gholami A, Emami SA, Mojarrab M. Cytotoxic and apoptotic effects of different extracts of Artemisia turanica Krasch. on K562 and HL-60 cell lines. *Sci World J.* 2013;2013. [[DOI:10.1155/2013/628073](https://doi.org/10.1155/2013/628073)] [[PMID](#)] [[PMCID](#)]
20. Afshari AR, Karimi Roshan M, Soukhtanloo M, et al. Cytotoxic effects of auraptene against a human malignant glioblastoma cell line. *Avicenna J Phytomed.* 2019;9(4):334.
21. Le L, Fu H, Lv Q, et al. The protective effects of the native flavanone flavanone on neuronal cells damaged by 6-OHDA. *Phytomedicine.* 2019;53:193-204. [[DOI:10.1016/j.phymed.2018.09.005](https://doi.org/10.1016/j.phymed.2018.09.005)] [[PMID](#)]
22. Andreeva-Gateva P, Traikov L, Sabit Z, et al. Antioxidant effect of alpha-lipoic acid in 6-hydroxydopamine unilateral intrastriatal injected rats. *Antioxidants.* 2020; 9(2):122. [[PMCID](#)] [[DOI:10.3390/antiox9020122](https://doi.org/10.3390/antiox9020122)] [[PMID](#)]
23. Kesh S, Kannan RR, Balakrishnan A. Naringenin alleviates 6-hydroxydopamine induced Parkinsonism in SHSY5Y cells and zebrafish model. *Comp Biochem Physiol Part - C: Toxicol Pharmacol.* 2021; 239:108893. [[DOI:10.1016/j.cbpc.2020.108893](https://doi.org/10.1016/j.cbpc.2020.108893)] [[PMID](#)]
24. Dirin MM, Mousavi SH, Afshari AR, Tabrizian K, Ashrafi MH. Potential drug-drug interactions in prescriptions dispensed in community and hospital pharmacies in East of Iran. *J Res Pharm*

- Pract. 2014;3(3):104. [PMID] [PMCID] [DOI:10.4103/2279-042X.141118]
25. Shafiee-Nick R, Afshari AR, Mousavi SH, et al. A comprehensive review on the potential therapeutic benefits of phosphodiesterase inhibitors on cardiovascular diseases. *Biomed Pharmacother.* 2017;94:541-56. [DOI:10.1016/j.biopha.2017.07.084] [PMID]
 26. Boroushaki MT, Mollazadeh H, Afshari AR. Pomegranate seed oil: A comprehensive review on its therapeutic effects. *Int J Pharm Sci Rev Res.* 2016;7(2):430.
 27. Mollazadeh H, Afshari AR, Hosseinzadeh H. Review on the potential therapeutic roles of nigella sativa in the treatment of patients with cancer: Involvement of apoptosis:-black cumin and cancer. *J Pharmacopunct.* 2017;20(3):158. [DOI:10.3831/KPLI.2017.20.019] [PMID] [PMCID]
 28. Afshari AR, Roshan MK, Soukhtanloo M, et al. Investigation of cytotoxic and apoptogenic effects of terminalia chebula hydro-alcoholic extract on glioblastoma cell line. *Shefaye Khatam.* 2018;5:14-23. [DOI:10.29252/shefa.6.4.14]
 29. Guo S, Bezard E, Zhao B. Protective effect of green tea polyphenols on the SH-SY5Y cells against 6-OHDA induced apoptosis through ROS-NO pathway. *Free Radic Biol Med.* 2005;39(5):682-95. [PMID] [DOI:10.1016/j.freeradbiomed.2005.04.022]
 30. Velikova V, Yordanov I, Edreva A. Oxidative stress and some antioxidant systems in acid rain-treated bean plants: protective role of exogenous polyamines. *Plant Sci.* 2000;151(1):59-66. [DOI:10.1016/S0168-9452(99)00197-1]
 31. Takase H, Liang AC, Miyamoto N, et al. Protective effects of a radical scavenger edaravone on oligodendrocyte precursor cells against oxidative stress. *Neurosci Lett.* 2018; 668:120-5. [DOI:10.1016/j.neulet.2018.01.018] [PMID] [PMCID]
 32. Quispe RL, Jaramillo ML, Galant LS, et al. Diphenyl diselenide protects neuronal cells against oxidative stress and mitochondrial dysfunction: Involvement of the glutathione-dependent antioxidant system. *Redox biology.* 2019;20:118-29. [PMCID] [PMID] [DOI:10.1016/j.redox.2018.09.014]
 33. Fang L, Ren D, Wang Z, Liu C, Wang J, Min W. Protective role of hazelnut peptides on oxidative stress injury in human umbilical vein endothelial cells. *J Food Biochem.* 2019;43(3):e12722. [DOI:10.1111/jfbc.12722] [PMID]
 34. Hosseini A, Sheikh S, Soukhtanloo M, et al. The effect of hydro-alcoholic extract of rheum turkestanicum roots against oxidative stress in endothelial cells. *Int J Prev Med.* 2020;11:122. [DOI:10.4103/ijpvm.IJPVM_386_19] [PMID] [PMCID]
 35. Crosti N, Servidei T, Bajer J, Serra A. Modification of the 6-hydroxydopamine technique for the correct determination of superoxide dismutase. *J Clin Chem Clin Biochem.* 1987;25(4):265-6.
 36. Perumal A, Gopal V, Tordzro W, Cooper T, Cadet J. Vitamin E attenuates the toxic effects of 6-hydroxydopamine on free radical scavenging systems in rat brain. *Brain Res Bull.* 1992;29(5): 699-701. [DOI:10.1016/0361-9230(92)90142-K]
 37. Baluchnejadmojarad T, Mansouri M, Ghalami J, Mokhtari Z, Roghani M. Sesamin imparts neuroprotection against intrastriatal 6-hydroxydopamine toxicity by inhibition of astroglial activation, apoptosis, and oxidative stress. *Biomed Pharmacother.* 2017;88:754-61. [DOI:10.1016/j.biopha.2017.01.123] [PMID]
 38. Rajabian A, Sadeghnia H-R, Moradzadeh M, Hosseini A. Rheum turkestanicum reduces glutamate toxicity in PC12 and N2a cell lines. *Folia neuropathol.* 2018;56(4):354-61. [DOI:10.5114/fn.2018.80869] [PMID]

How to Cite This Article:

Rashidi R, Afshari A R, Mollazadeh H, Hosseini A. Rheum turkestanicum Janisch Root Extract Mitigates 6-OHDA-Induced Neuronal Toxicity Against Human Neuroblastoma SH-SY5Y Cells. *J Adv Med Biomed Res.* 2022; 30(143): 553-60.

Download citation:

[BibTeX](#) | [RIS](#) | [EndNote](#) | [Medlars](#) | [ProCite](#) | [Reference Manager](#) | [RefWorks](#)

Send citation to:

 [Mendeley](#)  [Zotero](#)  [RefWorks](#) [RefWorks](#)