

## Comparison of RT-PCR and CT Scan in Terms of Their Diagnostic Performance for Coronavirus 2019 in Yazd, Iran

Atefeh Daya<sup>1</sup> , Nasim Namiranian<sup>2</sup> , Seyyed Kazem Razavi Ratki<sup>3</sup> ,  
Amir Pasha Amlshahbaz<sup>3</sup> , Roghaye Razavi<sup>2</sup> , Reza Nafisi Moghadam<sup>1,3\*</sup> 

1. Dept. of Radiology, Shahid Sadoughi Hospital, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
2. Yazd Diabetes Research Center, Shahid Sadoughi University of Medical Sciences, Yazd, Iran
3. Dept. of Radiology, Afshar Hospital, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

### Article Info

doi [10.30699/jambs.31.146.261](https://doi.org/10.30699/jambs.31.146.261)

Received: 2022/08/08;

Accepted: 2023/01/20;

Published Online: 26 June 2023;

Use your device to scan and read the article online



### Corresponding Information:

**Reza Nafisi-Moghadam,**

Dept. of Radiology, Shahid Sadoughi Hospital, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

E-Mail: [nafisi.moghadam@gmail.com](mailto:nafisi.moghadam@gmail.com)

### ABSTRACT

**Background & Objective:** Due to the importance of the rapid diagnosis of SARS-COV-2, many studies have been performed on various diagnostic methods. In this vein, the aim of this study is to compare the diagnostic value of chest CT with RT-PCR in patients with suspected COVID-19.

**Materials & Methods:** This cross-sectional research was conducted at Shahid Sadoughi Hospital in Yazd. A total of 531 patients were randomly referred to the CT scan department for chest imaging (spiral or HRCT). Based on the PCR findings, they were divided into: positive and negative PCR groups. The CT scan findings were then recorded in a data collection form. Finally, the CT scan results of the two groups were compared.

**Results:** In this study, 531 patients (306 males and 225 females with the mean age of  $55.14 \pm 19.7$ ) were examined. The findings of reverse halo ( $P = 0.000$ ) and strict consolidation ( $P = 0.001$ ) in CT scan were significantly different for the positive and negative PCR groups. Through the comparative analysis of the results, the sensitivity of CT scan and PCR findings emerged to be 97.42 and 55.75, respectively. However, the specificity of both diagnostic methods was 100%.

**Conclusion:** The results of this study show that, owing to the higher sensitivity of chest CT for the diagnosis of COVID-19 pneumonia, performing CT scan for quick diagnosis is recommended for COVID suspected people with negative RT-PCR test results.

**Keywords:** COVID-19, Pneumonia, Diagnosis, CT- Scan, RT- PCR



Copyright © 2023, This is an original open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permits copy and redistribution of the material just in noncommercial usages with proper citation.

## Introduction

Following the outbreak of an unknown type of pneumonia in Wuhan, China, in December 2019, the International Virus Classification Committee identified a new coronavirus from the RNA family of viruses called SARS-CoV-2, which can infect humans (1-3). Then, due to the spread of the disease worldwide, the WHO labeled that pneumonia as a pandemic that could threaten public health and the global economy and as a factor that could destabilize social communities (4-6). Since then, while the symptoms of the disease have been mild to moderate in most patients, some have developed the acute respiratory distress syndrome (ARDS), suffered from multifocal dysfunction, and died (7).

Because the virus can be transmitted rapidly from human to human, it is important to diagnose the disease in the early stages and isolate the infected individuals from the healthy population, thus reducing the transmission of

the infection to others in the society and preventing its prevalence (3, 8, 9). To this end, the existence of COVID-19 virus should be checked in patients with common clinical symptoms (e.g., fever, cough and hypoxemia) (10). The reverse transcription polymerase chain reaction (RT-PCR) is currently the reference standard with very high specificity to identify COVID-19 infection. According to recent reports, the sensitivity of RT-PCR for swab throat samples is limited; it is from 50% to 62%, which is too low for screening (11). So, using this method can leave many COVID-19 patients un-identified and, as a result, lead to the rapid transmission of the virus to others (9). Chest CT sensitivity for the diagnosis of COVID-19 has been reported to be high (from 60 to 98%) (12, 13). A previous study showed poor (25%) chest CT specificity for the diagnosis of COVID-19 (9). Overall, studies have shown that, compared to RTPCR, chest CT is of higher sensitivity for detecting COVID-19 (9). As a study found,

3% of the patients had a negative RT-PCR if their chest CT confirmed COVID-19 viral pneumonia (14). As shown in other studies, some patients with positive RT-PCR had normal chest CT. For example, Bernheim et al. reported normal chest CT in 56% of the patients with clinical symptoms (13, 15). In line with these studies, the aim of the present research is to compare the diagnostic values of chest CT and RT-PCR for the patients suspected of COVID-19.

## Materials and Methods

### Study design and participants

After the research was approved in the Ethics Committee of Shahid Sadoughi University of Medical Sciences of Yazd (IR.SSU.MEDICINE.REC.1399.131), the purpose of the study and the steps of the work were explained to all the participants, and written informed consent was obtained from them. This cross-sectional study was conducted from March 20, 2020 to June 20, 2020 on all those who had referred to Shahid Sadoughi Hospital in Yazd and were suspected of acute coronavirus 2 respiratory syndrome infection. The clinical symptoms were fever, cough, short breath, body aches and headaches. The patients who did not consent to CT scan after the procedure was explained to them or whose RT-PCR and CT data were not available were excluded from the study.

### Chest CT scan protocol and RT-PCR

Over a period of 24 hours or less, 531 patients suspected of COVID-19 underwent chest CT (spiral or HRCT, 1.2 mm incision in inhaling position, TOSHIBA medical systems, Otawara, Japan 16 slices / Alexion) and an RT-PCR test with throat samples. The sensitivity and specificity of the PCR test were calculated in a quadratic table, and the test emerged to be gold standard. The patients' demographic information and the CT scan findings were recorded in the forms designed by the radiologist. The patients were divided into one of the several groups based on the characteristics of their chest CT scan, including the existence of GGO, consolidation, the number of lobes involved, pleural effusion, nodules,

lymphadenopathy, and the existence of underlying lung diseases such as emphysema. Then, for the RT-PCR test, the participants were asked to refer to Shahid Sadoughi Medical Genetics Laboratory in Yazd in 24 hours or less. The samples were taken by Dacron swabs and transferred to a respiratory medium for RNA extraction. In the next step, a diagnostic test was performed by the RT-PCR method using a kit (Novel Coronavirus Diagnostic Kit, Sensure Biotech Inc. China). A radiologist who was unaware of the patients' RT-PCR results reviewed the CT scan findings and recorded them in special forms. Then, the RT-PCR data were collected from the electronic medical records based on PCR findings, and the patients were divided into positive and negative PCR groups. The diagnosis of COVID-19 infection was done based on the preliminary chest CT and RT-PCR findings.

### Statistical analysis:

The statistical analyze were performed using the SPSS 22 software. The continuous data were displayed as mean  $\pm$  standard deviation and the stratified variables as number (percentage). Then, a t-test and a chi-square test served to compare the two groups. In this regard,  $P < 0.05$  showed a statistically significant difference.

## Results

### Demographic characteristics

A total of 531 patients who were suspected of COVID-19 (306 males and 225 females with a mean age of  $55.14 \pm 19.7$ ) referred to Shahid Sadoughi Hospital in Yazd, were enrolled in the study according to the inclusion criteria, and underwent chest CT and RT-PCR for 24 hours. Nine patients were excluded from the study because the time interval between their chest CT and RT-PCR test was more than 24 hours, There was not access to the chest CT scans of some patients ( $N = 6$ ) because they were transferred to other hospitals; Thus, the RT-PCR findings of 522 patients and the chest CT of 506 of them were available for the final analysis. The participants' demographic characteristics are reported in [Table 1](#).

**Table 1. COVID-19 patient' demographic characteristics as well as initial RT- PCR and Chest CT imaging findings**

Variables	Values	
Age, Mean ( $\pm$ SD)	55.14 ( $\pm$ 19.7)	
<b>Sex, N (%)</b>		
Male	306 (57.6)	
Female	225 (46.4)	
<b>Results, N (%)</b>		
Initial RT- PCR assay	+	291 (55.7)
	-	231 (44.3)
<b>Pattern of the initial chest CT imaging, N (%)</b>		
GGO	+	467 (93.3)

Variables	Values
	- 39 (7.7)
	+ 297 (58.7)
<b>Patchy consolidation</b>	- 209 (41.3)
	+ 84 (16.6)
<b>Strict consolidation</b>	- 422 (83.4)
	+ 101 (19.9)
<b>Crazy paving</b>	- 405 (80.1)
	+ 75 (14.8)
<b>Reverse halo</b>	- 431 (85.2)
	+ 23 (4)
<b>Plural effusion</b>	- 583 (96)
	+ 2 (0.3)
<b>pneumothorax</b>	- 504 (99.7)
	+ 5 (0.9)
<b>emphysema</b>	- 501 (99.1)
<b>Distribution of the findings in the images, N (%)</b>	
<b>Central</b>	24 (4.8)
<b>Peripheral</b>	390 (77.2)
<b>Peripheral/central</b>	91 (18)

- RT-PCR: reverse transcription polymerase chain reaction; GGO: ground glass opacities

### CT Findings

As the computed tomography evaluation of the participants showed, the highest values belonged to the GGO and then patchy consolidation groups. These values were obtained through the CT scans of 467(93.3%) and 297(58.7%) patients. In addition, the findings on strict consolidation, crazy paving and reverse halo were recorded for 84 (16.6%), 101 (19.9%) and 75 (14.8%) patients suspected with COVID- 19, respectively. According to the results, the incidence of complications such as pleural effusion and pneumothorax was very low among the patients. Also, the study of the distribution of the lung lesions on the

CT scans of the participants showed that the most involvement was in the peripheral part of the lung ( $P = 0.002$  in 390 of 505 or 77.2% of the cases) (Table 2). Some COVID suspected patients with negative RT-PCR test results had chest CT findings on turbidity [266 of 286], GGO [166 of 286], and patchy consolidation and crazy paving [69 of 286], but the differences were not statistically significant. In terms of negative RT-PCR results and positive reverse halo findings, however, the participants were statistically different (43 of 286;  $P = 0.000$ ); Significant differences were also detected in the negative RT-PCR test results and the positive result on strict consolidation (62 of 286;  $P = 0.001$ ).

**Table 2.** Comparison of the participants in terms of RT- PCR, chest CT imaging and distribution of lesions

Parameters	RT- PCR				P- value	
	-		+			
	Male	Female	Male	Female		
<b>GGO</b>	-	6 (1.2)	14 (2.8)	11 (2.2)	6 (1.2)	0.782
	+	111 (22.2)	155 (31.1)	90 (18)	105 (21)	
<b>Patchy consolidation</b>	-	56 (11.2)	64 (12.8)	41 (8.2)	44 (8.8)	0.183
	+	61 (12.2)	105 (21)	60 (12)	67 (13.4)	

Parameters	RT- PCR				P- value	
	-		+			
	Male	Female	Male	Female		
Reverse halo	-	112 (22.4)	131 (26.3)	89 (17.8)	91 (18.2)	0.000
	+	5 (1)	38 (7.6)	12 (2.4)	20 (4)	
Strict consolidation	-	102 (20.4)	122 (24.4)	93 (18.6)	97 (19.4)	0.001
	+	15 (3)	47 (9.4)	8 (1.6)	14 (2.8)	
Crazy paving	-	91 (18.2)	126 (25.3)	93 (18.6)	91 (18.2)	0.052
	+	26 (5.2)	43 (8.6)	8 (1.6)	20 (4)	
Central		10 (2)	4 (0.8)	8 (1.6)	1 (0.2)	
Peripheral		90 (18.1)	135 (27.1)	74 (14.8)	85 (17.1)	0.002
Peripheral/ Central		17 (3.4)	30 (6)	18 (3.6)	25 (5)	

- RT-PCR: reverse transcription polymerase chain reaction; GGO: ground glass opacities

### Comparative diagnostic results

At the beginning of the study, the RT-PCR test results were positive in 291 patients, abnormal CT scan findings were observed in 506 (95.3%) participants, and the chest CT results for COVID-19 were positive in 218 (94.4%) of 231 patients whose RT-PCR result was negative. The diagnostic data obtained from the primary RT-PCR tests and CT scans are reported in [Table 3](#). According to the comparative analysis of the CT scan findings and the initial RT-PCR results, the sensitivity and accuracy of COVID-19 detection with the initial RT-PCR tests was 55.75%, while it was

97.42% with the initial CT scan as for specificity, it was found to be 100% in both initial diagnostic modes. Generally, there was no statistically significant difference between the initial CT detection rate and the initial RT-PCR ( $P = 0.317$ ).

Sensitivity

$$= \frac{\text{number of true positives}}{\text{number of true positives} + \text{number of false negatives}}$$

Specificity

$$= \frac{\text{number of true negatives}}{\text{number of true negatives} + \text{number of false positives}}$$

**Table 3. Diagnostic rates of RT-PCR and CT scan**

Diagnostic rate (%)	Sensitivity	Specificity	Accuracy
Initial RT- PCR	55.75	100	55.75
Initial CT imaging	97.42	100	97.42
P- value	0.0001	1.000	0.0001

## Discussion

According to the literature on COVID-19, the most important measure to control and prevent the spread of the disease is the rapid diagnosis of the infected individuals, their rapid isolation and quarantine, and the identification and isolation of all those in close contact with them from the healthy population (16). In this regard, RT-PCR is of relatively low diagnostic sensitivity, has limited diagnostic kits, and is practiced with insufficient skills in many countries. Indeed, it fails to identify many people infected with COVID-19, which leads to further transmission of the virus (11). Therefore, this method can-not be the only reliable tool for COVID-19 screening (11).

In this study, the PCR test was positive for 291 patients infected with COVID-19, but the chest CT of

many participants showed a variety of findings on variables such as GGO, patchy consolidation, and peripheral distribution of lesions. As expected, the most common of these findings was GGO, and the rarest were pleural effusion and pneumothorax. According to the statistical data, the CT scans of the positive and negative PCR groups were significantly different only in terms of reverse halo data and lesion distribution location. However, GGO lesions and patchy consolidation were not significantly different in the two groups.

The diagnostic sensitivity of the RT-PCR test in this study was 55.75%; the results here are in line with the findings of some other studies in this field, such as Ai et al. (9). In a study of 82 hospitalized patients

conducted by He et al., the sensitivity and specificity of RT-PCR was found to be 79% and 100%, respectively (11). This difference can be due to some factors such as the differences in the distribution of the kits used, the method and quality of testing, and the viral load on the respiratory tract (17).

In the present study, the diagnostic sensitivity of the chest CT test conducted on COVID-19 patients was 97.42%. In a similar study on 36 COVID-19 pneumonia- suspected patients conducted by Long et al. the CT sensitivity of the chest emerged to be 97.2%, which confirms the CT scan findings obtained in this study. In terms of the PCR sensitivity, however, Long et al. (ibid) obtained a value of 83.3%, which is higher than that in the present study (3). The sensitivity of imaging calculated by Hao et al. is consistent with the results of our investigation (18). Also, Dangis et al. evaluated the diagnostic sensitivity, specificity and accuracy of the CT scans of 192 COVID-19 patients. Those factors were 86.7%, 93.6% and 91.1% respectively. In another study on symptomatic patients for 48 hours, the sensitivity, specificity and accuracy of the CT scans were 95.6%, 93.2% and 91.5%, respectively (19). Moreover, in a study in 2020, the sensitivity and specificity of chest CT were reported to be 77% and 96%, respectively (11). In a large study on 919 patients in Wuhan, the chest CT sensitivity was 97% (9). Generally, the chest CT sensitivity in studies for the diagnosis of COVID-19 has been up to 98% (20). This suggests that chest CT imaging, due to its high sensitivity, easy access and cost-effectiveness, can serve as an effective diagnostic method to separate COVID-19 suspected people from others (11).

The findings of this study point to a statistically significant difference between chest CT and RT-PCR in terms of sensitivity. Fang et al. demonstrated that chest CT was significantly more sensitive than RT-PCR (98% and 71%, respectively) (20), which is in line with the findings of our study. In the study conducted by He et al., the difference of the two techniques was not significant (11). In the same vein of research, Bai et al. found no significant relationship between rt-PCR and chest in terms of sensitivity and specificity (21).

In a study in Tabriz, Portahmasebi et al. concluded that CT scan is suitable for assessing the disease and its severity, but it is not recommended for screening COVID-19 suspected ones because of the harmful ionizing radiation used in it (8). In this respect, some studies have shown that at limited dose of radiation in chest CT does not cause cancer (22); Anyway, it is suggested that this method be used with caution for certain age groups and vulnerable people.

In an extensive study by Yang et al., several pieces of research with various statistical populations were reviewed and analyzed for their findings, which is worth considering. It was found that, in many cases, normal CT scans had been done on the early days of the disease despite positive PCR test results. This may be due to the fact that COVID-19 infected people in areas

outside the epidemic have only the initial and mild symptoms of the disease. As a result, there is no pneumonia or lung involvement in these cases, and their chest CT scan is normal (11, 14). Herein, it is suggested that further studies be designed with larger size samples, over longer periods of time, in several medical centers and by using gold standard diagnostic methods to yield more reliable and helpful results.

## Conclusion

As shown by many studies in the short time since the onset of the COVID-19 pandemic, the early diagnosis of the disease and, consequently, the disruption of its transmission chain are among the major concerns of all medical systems in the world although there are many controversies over procedures and findings. Overall, the results of this study suggest that RT-PCR may be negative in many COVID-19-infected individuals, while chest CT scans are more sensitive for the detection of the virus as well as faster and more accurate for the evaluation of those patients. Hence, due to the higher sensitivity of chest CT for the diagnosis of COVID-19 pneumonia, it is recommended that people who are clinically suspected of COVID-19 but have a negative RT-PCR test be evaluated for a quick diagnosis of infection using a chest CT scan. If a person proves to be suspected of COVID-19 pneumonia on a chest CT scan, it is necessary to isolate him or her from other people in the society and to start appropriate treatment to prevent further infection or outbreak of the disease.

## Acknowledgments

The authors thank all the study participants and the hospital staff for their support and cooperation in this study.

## Funding

This research was financially supported by the Vice Chancellor of Research at Shahid Sadoughi University of Medical Sciences and Health Services in Yazd.

## Conflict of Interest

The authors declare no conflict of interests regarding this study.

## References

1. Phelan AL, Katz R, Gostin LO. The novel coronavirus originating in Wuhan, China: challenges for global health governance. *Jama*. 2020;323(8):709-10. [DOI:10.1001/jama.2020.1097] [PMID]
2. Wu Y, Ho W, Huang Y, et al. SARS-CoV-2 is an appropriate name for the new coronavirus. *The*

- Lancet. 2020;395(10228):949-50. [DOI:10.1016/S0140-6736(20)30557-2] [PMID]
3. Long C, Xu H, Shen Q, et al. Diagnosis of the coronavirus disease (COVID-19): rRT-PCR or CT? *Europ J Radiol.* 2020;126:108961. [PMID] [DOI:10.1016/j.ejrad.2020.108961] [PMCID]
  4. Legido-Quigley H, Asgari N, Teo YY, et al. Are high-performing health systems resilient against the COVID-19 epidemic? *The Lancet.* 2020; 395(10227):848-50. [PMID] [DOI:10.1016/S0140-6736(20)30551-1]
  5. Wang P, Anderson N, Pan Y, et al. The SARS-CoV-2 outbreak: diagnosis, infection prevention, and public perception. *Clin Chem.* 2020;66(5): 644-51. [DOI:10.1093/clinchem/hvaa080] [PMID] [PMCID]
  6. Khot WY, Nadkar MY. The 2019 novel coronavirus outbreak-a global threat. *J Assoc Physicians India.* 2020;68(3):67-71.
  7. Ye Q, Wang B, Mao J. The pathogenesis and treatment of the Cytokine Storm'in COVID-19. *J Infect.* 2020;80(6):607-13. [PMID] [PMCID] [DOI:10.1016/j.jinf.2020.03.037]
  8. Poortahmasebi V, Zandi M, Soltani S, Jazayeri SM. Clinical performance of RT-PCR and chest CT scan for COVID-19 diagnosis; a systematic review. *Front Emerg Med.* 2020;4(2s):e57-e.
  9. Ai T, Yang Z, Hou H, et al. Correlation of chest CT and RT-PCR testing in coronavirus disease 2019 (COVID-19) in China: a report of 1014 cases. *Radiology.* 2020; 296(2):E32-E40. [DOI:10.1148/radiol.2020200642] [PMID] [PMCID]
  10. Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *Jama.* 2020;323(11):1061-9. [DOI:10.1001/jama.2020.1585] [PMID] [PMCID]
  11. He JL, Luo L, Luo ZD, et al. Diagnostic performance between CT and initial real-time RT-PCR for clinically suspected 2019 coronavirus disease (COVID-19) patients outside Wuhan, China. *Respir Med.* 2020;168:105980. [DOI:10.1016/j.rmed.2020.105980] [PMID] [PMCID]
  12. Ng MY, Lee EY, Yang J, et al. Imaging profile of the COVID-19 infection: radiologic findings and literature review. *Radiol Cardiothorac Imaging.* 2020;2(1):e200034. [PMCID] [DOI:10.1148/ryct.2020200034] [PMID]
  13. Bernheim A, Mei X, Huang M, et al. Chest CT findings in coronavirus disease-19 (COVID-19): relationship to duration of infection. *Radiology.* 2020. [DOI:10.1148/radiol.2020200463] [PMID] [PMCID]
  14. Xie X, Zhong Z, Zhao W, Zheng C, Wang F, Liu J. Chest CT for typical 2019-nCoV pneumonia: relationship to negative RT-PCR testing. *Radiology.* 2020.
  15. Yang W, Yan F. Patients with RT-PCR-confirmed COVID-19 and normal chest CT. *Radiology.* 2020;295(2):E3-E. [PMID] [PMCID] [DOI:10.1148/radiol.2020200702]
  16. Zhang Z, Yao W, Wang Y, Long C, Fu X. Wuhan and Hubei COVID-19 mortality analysis reveals the critical role of timely supply of medical resources. *J Infect.* 2020;81(1):147-78. [DOI:10.1016/j.jinf.2020.03.018]
  17. Chan JF-W, Yip CC-Y, To KK-W, et al. Improved molecular diagnosis of COVID-19 by the novel, highly sensitive and specific COVID-19-RdRp/Hel real-time reverse transcription-PCR assay validated in vitro and with clinical specimens. *J Clin Microbiol.* 2020;58(5):e00310-20. [DOI:10.1128/JCM.00310-20] [PMID] [PMCID]
  18. Hao W, Li M. Clinical diagnostic value of CT imaging in COVID-19 with multiple negative RT-PCR testing. *Travel Med Infect Dis.* 2020;34: 101627. [DOI:10.1016/j.tmaid.2020.101627] [PMID] [PMCID]
  19. Dangis A, Gieraerts C, De Bruecker Y, et al. Accuracy and reproducibility of low-dose submillisievert chest CT for the diagnosis of COVID-19. *Radiol Cardiothorac Imaging.* 2020; 2(2):e200196. [PMID] [PMCID] [DOI:10.1148/ryct.2020200196]
  20. Fang Y, Zhang H, Xie J, et al. Sensitivity of chest CT for COVID-19: comparison to RT-PCR. *Radiology.* 2020. [PMID] [PMCID] [DOI:10.1148/radiol.2020200432]
  21. Bai HX, Hsieh B, Xiong Z, et al. Performance of radiologists in differentiating COVID-19 from non-COVID-19 viral pneumonia at chest CT. *Radiology.* 2020;296(2):E46-E54. [PMCID] [DOI:10.1148/radiol.2020200823] [PMID]
  22. Schultz CH, Fairley R, Murphy LS L, Doss M. The risk of cancer from CT scans and other sources of low-dose radiation: a critical appraisal of methodologic quality. *Prehosp Disaster Med.* 2020;35(1):3-16. [DOI:10.1017/S1049023X1900520X] [PMID]

**How to Cite This Article:**

Daya A, Namiranian N, Razavi Ratki S K, Amlshahbaz A P, Razavi R, Nafisi Moghadam R. Comparison of RT-PCR and CT Scan in Terms of Their Diagnostic Performance for Coronavirus 2019 in Yazd, Iran. *J Adv Med Biomed Res.* 2023; 31(146): 261-7.

**Download citation:**

[BibTeX](#) | [RIS](#) | [EndNote](#) | [Medlars](#) | [ProCite](#) | [Reference Manager](#) | [RefWorks](#)

**Send citation to:**

 [Mendeley](#)  [Zotero](#)  [RefWorks](#) [RefWorks](#)