

# Antiepileptic Role of $\beta$ -alanine on Pentylenetetrazol-Induced Convulsions during Estrus Cycle in Rat

Shahin Hassanzadeh <sup>1\*</sup> , Sahar Hajimoradi <sup>1</sup> , Faezeh Malekpour <sup>2</sup> 

1. Division of Physiology, Department of Basic Sciences, Faculty of Veterinary Medicine, Science and Research

Branch, Islamic Azad University, Tehran, Iran

2. Iran University of Medical Sciences, Tehran, Iran

## Article Info

 [10.30699/jambr.32.154.388](https://doi.org/10.30699/jambr.32.154.388)

Received: 2023/10/31;

Accepted: 2024/11/31;

Published Online: 31 Dec 2024;

## ABSTRACT

**Background & Objective:** Epilepsy is one of the main neurological disorders and  $\beta$ -alanine has several beneficial in the nervous system. This study aimed to determine the antiepileptic effect of the  $\beta$ -alanine during estrous cycle in pentylenetetrazol (PTZ)-induced seizures in rat.

**Materials & Methods:** Thirty female rats were allocated into five experimental groups as control, sodium valproate (75 mg/kg), and  $\beta$ -alanine (15, 30 and 45 mg/kg) during estrus cycle (proestrus, estrus, metestrus and diestrus). Puberty was determined by vaginal smears whereby animals with two normal cycles were chosen. After administration of the sodium valproate or  $\beta$ -alanine, PTZ injection (80 mg/kg, i.p) was done. Animals were monitored for 30 minutes with initiation time of myoclonic seizures (ITMS), initiation time of tonic-clonic seizures (ITTS), and seizures' duration (SD) registered.

**Results:** Based on the findings,  $\beta$ -alanine (30 and 45 mg/kg) significantly increased the onset time of ITMS and ITTS compared to the control group ( $P<0.05$ ).  $\beta$ -alanine (30 and 45 mg/kg) significantly reduced SD compared to the control group ( $P<0.05$ ). The antiepileptic effect of  $\beta$ -alanine was more prominent during metestrus and diestrus than proestrus and estrus ( $P<0.05$ ).

**Conclusion:** It seems  $\beta$ -alanine has antiepileptic effects which are more prominent during proestrus and estrus than metestrus and diestrus.

**Keywords:** Antiepileptic,  $\beta$ -alanine, PTZ, Estrus, Rat

## Corresponding Information:

Shahin Hassanzadeh,  
Division of Physiology, Department of  
Basic Sciences, Faculty of Veterinary  
Medicine, Science and Research  
Branch, Islamic Azad University, Tehran, Iran  
E-Mail: [s.hassanzadeh@srbiau.ac.ir](mailto:s.hassanzadeh@srbiau.ac.ir)



Copyright © 2023, This is an original open-access article distributed under the terms of the Creative Commons Attribution-noncommercial 4.0 International License which permits copy and redistribution of the material just in noncommercial usages with proper citation.

## Introduction

The basic pathophysiology of epilepsy is not fully provoked but imbalance between excitatory and inhibitory anticonvulsants play important role (1). Several models of focal epilepsy are based on impairment of cortical GABAergic neurons and most anti-epileptic drugs act by inhibition of  $\text{Na}^+$  or  $\text{Ca}^{2+}$  channels (2). PTZ is commonly used for inducing general seizure in experimental catamenial epilepsy which acts by inhibition of the GABA<sub>A</sub> receptor (3). Catamenial epilepsy is a type of the seizure which happens in results of the changes in sexual hormone in women. There is ample evidence that ovarian estrogens contribute to the development of epilepsy, whereas progesterone has a protective effect against seizures

(4). Cyclic changes in ovarian hormones during estrous cycle may play in incidence of the catamenial epilepsy. Progesterone or its derivatives allopregnanolone and pregnanolone affect GABA receptors and reduce the occurrence of seizures (5). Enzyme-modulating antiepileptic drugs such as carbamazepine, phenobarbital sodium valproate is prescribed for epilepsy but they can alter the serum level of steroid hormones (6).

$\beta$ -alanine is the simplest  $\beta$ -amino acid within the human central nervous system as a neuromodulator (7).  $\beta$ -alanine has high affinity for normal and reactive astrocytes and competes with GABA for glial uptake (7).  $\beta$ -alanine interacts with no less than five

acknowledged receptor sites on NMDA and glycine receptor and GABA<sub>A</sub>, and also inhibits the glial GABA uptake mediated by GAT protein. Additionally,  $\beta$ -alanine is able to penetrate the blood brain barrier (7). Oral supplementation of the  $\beta$ -alanine improves exercise performance, inhibition of tumor growth and enhanced glucose sensitivity in patients with Parkinson's disease (9). Despite the fact that  $\beta$ -alanine play role of the in GABA<sub>A</sub>, and involvement of this receptors in catamenial epilepsy, this study aimed to determine antiepileptic effect of the  $\beta$ -alanine during estrous cycle in PTZ-induced seizures in rat.

## Materials and Methods

The sample consisted of thirty female Wistar rats weighing  $200 \pm 50$  g, which were categorized into 5 different experimental groups based on their estrus cycle phase (proestrus, estrus, metestrus, and diestrus). Animals were kept in accordance with European community regulations for laboratory animals, with standard conditions of  $22 \pm 1^\circ\text{C}$  and a 12-hour dark/light cycle. They had free access to food and water. Puberty was studied by analyzing vaginal smears. Rats with two normal estrous cycles were selected for the study and estrus synchronization was performed on them (10). Daily vaginal smears were used to determine the stage of estrous cycles by analyzing the most common cell type (11,12).

### Study procedure

In control group rat received saline while in experiment 2, animals received VPA (75 mg/kg, Sigma CAS Number, 1069-66-5). In experiments 3-5, animals received  $\beta$ -alanine (15, 30 and 45 mg/kg, Sigma CAS Number, 107-95-9), respectively. Then, animals received i.p administration of PTZ (80 mg /kg). Each experiment done during estrus cycle (proestrus, estrus, metestrus, and diestrus). Following the seizure being triggered, the animals' conduct was observed for a period of 30 minutes in order to evaluate the time it took for myoclonic seizures (ITMS) and tonic-clonic seizures (ITTS) to begin, as well as their duration (SD). Experiments were carried out between 9 and 12 in the

morning to minimize the influence of circadian rhythm on seizure vulnerability.

### Statistical Analysis


Data was analyzed by SPSS version 21 through a one-way analysis of variances (ANOVA) and Tukey multiple comparison tests, with results presented as mean  $\pm$  SD ( $P < 0.05$ ).

## Results


The effects of VPA,  $\beta$ -alanine (15, 30, and 45 mg/kg) on the ITMS is presented in figure 1. As observed, valproic acid injection significantly increased ITMS in the stages of proestrus, estrus, metestrus and diestrus compared to the control group ( $P < 0.05$ ).  $\beta$ -alanine (15 mg/kg) had no significant effect on ITMS in proestrus, estrus, metestrus and diestrus compared to control group ( $P > 0.05$ ).  $\beta$ -alanine (30, and 45 mg/kg) significantly inhibited the ITMS in the stages of proestrus, estrus, metestrus and diestrus in comparison to control group ( $P < 0.05$ ). The effects of the  $\beta$ -alanine were more prominent during metestrus and diestrus than proestrus, estrus ( $P < 0.05$ ).

As seen in figure 2, VPA injection significantly increased ITTS in proestrus, estrus, metestrus and diestrus stages compared to the control group ( $P < 0.05$ ).  $\beta$ -alanine (15 mg/kg) had no significant effect on ITTS in proestrus, estrus, metestrus and diestrus ( $P > 0.05$ ).  $\beta$ -alanine (30, and 45 mg/kg) significantly decreased ITTS in the stages of proestrus, estrus, metestrus and diestrus ( $P < 0.05$ ).

According to figure 3, VPA (75 mg/kg) significantly decreased SD in proestrus, estrus, metestrus and diestrus phases compared to the control group ( $P < 0.05$ ).  $\beta$ -alanine (15 mg/kg) had no significant effect on SD in proestrus, estrus, metestrus and diestrus stages ( $P > 0.05$ ).  $\beta$ -alanine (30, and 45 mg/kg) significantly decreased SD compared to the control group ( $P < 0.05$ ). The effects of the  $\beta$ -alanine were more prominent during metestrus and diestrus than proestrus, estrus ( $P < 0.05$ ).




**Figure 1.** Antiepileptic effects of  $\beta$ -alanine (15, 30, and 45 mg/kg) on the initiation time of myoclonic seizures (ITMS) (sec) during various phases of the estrous cycle. \*Asterisks indicate a significant difference in each estrous cycle phase compared with the control group ( $P < 0.05$ ). Different letters (a and b) indicate significant differences for each group in each estrous cycle phase ( $P < 0.05$ ). Data are presented as mean  $\pm$  SE.



**Figure 2.** Antiepileptic effects of  $\beta$ -alanine (15, 30, and 45 mg/kg) on the initiation time of tonic-clonic seizures (ITTS) (sec) during various phases of the estrous cycle. \*Asterisks show a significant difference in each estrous cycle phase

compared with the control group ( $P < 0.05$ ). Different letters (a, b, and c) display significant differences for each group in each estrous cycle phase ( $P < 0.05$ ). Data are presented as mean  $\pm$  SE.



**Figure 3.** Antiepileptic effects of  $\beta$ -alanine (15, 30, and 45 mg/kg) on seizure duration (sec) during various estrous cycle phases. \*Asterisks reveal a significant difference in each estrous cycle phase compared with the control group ( $P < 0.05$ ). Different letters (a and b) indicate significant differences for each group in each estrous cycle phase ( $P < 0.05$ ). Data are presented as mean  $\pm$  SE.

## Discussion

In epileptic women, fluctuations in ovarian steroids and their metabolites are responsible for catamenial epilepsy. Seizure incidence increases during the follicular phase and declines during the luteal phase (13). Elevated estrogen levels in the amygdala and hippocampus enhances seizure threshold while progesterone lowers it. Diminished progesterone levels in preovulatory phase leads to increased seizure incidence (14). Allopregnanolone is an allosteric metabolite of the progesterone and acts as modulator of GABA<sub>A</sub> receptors (15). Based on the main finding of the current study,  $\beta$ -alanine (30 and 45 mg/kg) increased the onset time of ITMS and ITTS and decreased SD. The antiepileptic effect of  $\beta$ -alanine was more prominent during metestrus and diestrus than proestrus and estrus.  $\beta$ -alanine can cross the blood brain barrier and serve as a marker for glial GABA uptake (16). Neuronal-glial studies have revealed  $\beta$ -alanine has a high affinity for glial GABA<sub>A</sub> receptors (17).

$\beta$ -Alanine is easily transported in the CNS, muscles, and other organs. In the brain  $\beta$ -alanine transporters by

GABA transporter (GAT). The GAT-1 and GAT-4 are found in the brain while GAT-2 and GAT-3 are found in the brain and peripheral tissues. GAT transports  $\beta$ -alanine, and  $\beta$ -alanine inhibits GABA uptake by GAT-2 (7). Despite direct mechanism for how  $\beta$ -alanine regulates GABA uptake is not fully elicited but it seems  $\text{Na}^+$  ions are required for  $\beta$ -alanine and GABA.  $\beta$ -alanine uptake is dependent on the concentration of both  $\text{Na}^+$  and  $\text{Cl}^-$  ions (18).

GABA is the primary inhibitory neurotransmitter in the brain, and its transmission plays a role in controlling mood, memory, and eating habits. Allopregnanolone, a progesterone metabolite, is a powerful modulator of GABA receptors with positive effects. Exposure to steroids that modulate GABA receptors or increased sensitivity to allopregnanolone can lead to hyperactivity or heightened GABAergic tone, and is linked to various symptoms as well as disorders (19). Administration of steroid antagonists that modulate GABA<sub>A</sub> receptors lowers the intensity of symptoms and conditions (20). The binding of GABA to its receptor leads to an increase in chloride

ion flow, membrane hyperpolarization, and reduction in neuronal firing rates (21-22). Direct GABA antagonists binding to GABAA receptors can lead to increased neuronal excitability and seizures while also interfering with GABA binding sites (23-25). A potential use of  $\beta$ -alanine, instead of carnosine, in aquaculture has been suggested to enhance the growth and feeding rate of mussels.  $\beta$ -alanine injections disrupted metabolites related to lipid metabolism. Specifically, metabolites involved in linoleic acid metabolism and steroid biosynthesis increased after  $\beta$ -alanine injection (26).

## Conclusion

In conclusion,  $\beta$ -alanine seems to have an antiepileptic role which is more prominent during metestrus and diestrus than proestrus and estrus. However, further research is required to determine accuracy of these findings in human trials.

## Acknowledgments

## References

- Chen K, Stieger KC, Kozai TD. Challenges and opportunities of advanced gliomodulation technologies for excitation-inhibition balance of brain networks. *Curr Opin Biotechnol.* 2021; 72:112-20. <https://doi.org/10.1016/j.copbio.2021.10.008> PMid:34773740 PMCid:PMC8671375
- Marafiga JR, Pasquetti MV, Calcagnotto ME. GABAergic interneurons in epilepsy: More than a simple change in inhibition. *Epilep Behav.* 2021; 121: 106935. <https://doi.org/10.1016/j.yebeh.2020.106935> PMid:32035792
- Pahwa P, Singh T, Goel RK. Anticonvulsant effect of Asparagus racemosus willd. In a mouse model of catamenial epilepsy. *Neurochem Res.* 2022;1-2. <https://doi.org/10.1007/s11064-021-03455-2> PMid:34623565
- Reddy DS, Thompson W, Calderara G. Molecular mechanisms of sex differences in epilepsy and seizure susceptibility in chemical, genetic and acquired epileptogenesis. *Neurosci Lett.* 2021; 750:135753. <https://doi.org/10.1016/j.neulet.2021.135753> PMid:33610673 PMCid:PMC7994197
- Miziak B, Chrościńska-Krawczyk M, Czuczwarcz SJ. Neurosteroids and seizure activity. *Frontiers in endocrinology.* 2020; 11:541802. <https://doi.org/10.3389/fendo.2020.541802> PMid:33117274 PMCid:PMC7561372
- Stephen LJ, Harden C, Tomson T, Brodie MJ. Management of epilepsy in women. *Lancet Neurol.* 2019; 18(5):481-91. [https://doi.org/10.1016/S1474-4422\(18\)30495-2](https://doi.org/10.1016/S1474-4422(18)30495-2) PMid:30857949
- Tiedje KE, Stevens K, Barnes S, Weaver DF.  $\beta$ -Alanine as a small molecule neurotransmitter. *Neurochem Int.* 2010; 57(3):177-88. <https://doi.org/10.1016/j.neuint.2010.06.001> PMid:20540981
- Dolan E, Swinton PA, Painelli VD, Stephens Hemingway B, Mazzolani B, Infante Smaira F, et al. A systematic risk assessment and meta-analysis on the use of oral  $\beta$ -alanine supplementation. *Adv Nutr.* 2019; 10(3):452-63. <https://doi.org/10.1093/advances/nmy115> PMid:30980076 PMCid:PMC6520041
- Zimmermann M. Ethical guidelines for investigations of experimental pain in conscious animals. *Pain.* 1983; 16: 109-110.

None.

## Authors' Contribution

## Conflict of Interest

The authors declare that they have no conflict of interest.

## Funding

None.

## Ethics Approval and consent to participate

The research protocol received approval from the Animal Ethics Committee at the Science and Research Branch of Islamic Azad University in Tehran, Iran (IR.IAU.SRB.REC.1401.597; 2023-2-16).

[https://doi.org/10.1016/0304-3959\(83\)90201-4](https://doi.org/10.1016/0304-3959(83)90201-4)  
PMid:6877845

10. Mattson RH, Cramer JA. Epilepsy, sex hormones, and antiepileptic drugs. *Epilepsia*. 1985; 26:S40-51.  
<https://doi.org/10.1111/j.1528-1157.1985.tb05723.x>  
PMid:3158512

11. Kaboutari J, Zendehdel M, Habibian S, Azimi M, Shaker M, Karimi B. The antiepileptic effect of sodium valproate during different phases of the estrous cycle in PTZ-induced seizures in rats. *J Physiol Biochem*. 2012; 68:155-61.  
<https://doi.org/10.1007/s13105-011-0127-x>  
PMid:22076911

12. Gutierrez EG, Crone NE, Kang JY, Carmenate YI, Krauss GL. Strategies for non-EEG seizure detection and timing for alerting and interventions with tonic-clonic seizures. *Epilepsia*. 2018; 59:36-41.  
<https://doi.org/10.1111/epi.14046>  
PMid:29873833

13. Zendehdel M, Kaboutari J, Ghadimi D, Hassanpour S. The antiepileptic effect of ghrelin during different phases of the estrous cycle in PTZ-induced seizures in rat. *Int J Peptide Res Therap*. 2014; 20:511-7.  
<https://doi.org/10.1007/s10989-014-9418-8>

14. Maguire MJ, Nevitt SJ. Treatments for seizures in catamenial (menstrual-related) epilepsy. *Coch Database System Rev*. 2019(10).  
<https://doi.org/10.1002/14651858.CD013225.pub2>  
PMid:31608992 PMCid:PMC6953347

15. Akk G, Germann AL, Sugasawa Y, Pierce SR, Evers AS, Steinbach JH. Enhancement of muscimol binding and gating by allosteric modulators of the GABAA receptor: relating occupancy to state functions. *Molec Pharmacol*. 2020; 98(4):303-13.  
<https://doi.org/10.1124/molpharm.120.000066>  
PMid:32873746

16. Al-Khawaja A, Petersen JG, Damgaard M, Jensen MH, Vogensen SB, Lie ME, et al. Pharmacological identification of a guanidine-containing  $\beta$ -alanine analogue with low micromolar potency and selectivity for the betaine/GABA transporter 1 (BGT1). *Neurochem Res*. 2014; 39:1988-96.  
<https://doi.org/10.1007/s11064-014-1336-9>  
PMid:24852577

17. Robitaille Y, Kemball K, Sherwin AL.  $\beta$ -Alanine uptake is upregulated in FeC13-induced cortical scars. *J Neurol Sci*. 1995; 134:95-101.

[https://doi.org/10.1016/0022-510X\(95\)00214-M](https://doi.org/10.1016/0022-510X(95)00214-M)  
PMid:8847551

18. Wang Z, Xie R, Yang X, Yin H, Li X, Liu T, et al. Female mice lacking ER $\beta$  display excitatory/inhibitory synaptic imbalance to drive the pathogenesis of temporal lobe epilepsy. *Theranostics*. 2021; 11(12):6074-6089.  
<https://doi.org/10.7150/thno.56331>  
PMid:33897900 PMCid:PMC8058727

19. Zanboori A, Tamaddonfar E, Mojtabahedin A. The effect of intracerebroventricular injection of histamine in visceral nociception induced by acetic acid in rats. *Indian J Pharmacol*. 2010;42(5):289.  
<https://doi.org/10.4103/0253-7613.70157>  
PMid:21206620 PMCid:PMC2959211

20. Bäckström T, Das R, Bixo M. Positive GABAA receptor modulating steroids and their antagonists: Implications for clinical treatments. *J Neuroendocrinol*. 2022; 34(2):e13013.  
<https://doi.org/10.1111/jne.13013>  
PMid:34337790

21. Wanleenuwat P, Suntharampillai N, Iwanowski P. Antibiotic-induced epileptic seizures: mechanisms of action and clinical considerations. *Seizure*. 2020; 81:167-74.  
<https://doi.org/10.1016/j.seizure.2020.08.012>  
PMid:32827980

22. Mayor Jr F, Marvizon JG, Aragon MC, Gimenez C, Valdivieso F. Glycine transport into plasma-membrane vesicles derived from rat brain synaptosomes. *Biochem J*. 1981; 198(3):535-41.  
<https://doi.org/10.1042/bj1980535>  
PMid:7326021 PMCid:PMC1163299

23. Stephen LJ, Harden C, Tomson T, Brodie MJ. Management of epilepsy in women. *Lancet Neurol*. 2019; 18(5):481-91.  
[https://doi.org/10.1016/S1474-4422\(18\)30495-2](https://doi.org/10.1016/S1474-4422(18)30495-2)  
PMid:30857949

24. Tinelli C, Di Pino A, Ficulle E, Marcelli S, Feligioni M. Hyperhomocysteinemia as a risk factor and potential nutraceutical target for certain pathologies. *Front Nutr*. 2019; 6:49.  
<https://doi.org/10.3389/fnut.2019.00049>  
PMid:31069230 PMCid:PMC6491750

25. Van Luijtelaar G, Budziszewska B, Jaworska-Feil L, et al. The ovarian hormones and absence epilepsy: a long term EEG study and pharmacological effects in a genetic absence model. *Epilepsy Res*. 2001; 46:225-239.

[https://doi.org/10.1016/S0920-1211\(01\)00277-7](https://doi.org/10.1016/S0920-1211(01)00277-7)  
PMid:11518624

26. Wang CY, Yan XJ, He JY, Buttino I, Pan C, Fan MH, Guo BY, Zhang XL, Liao Z. Responses to  $\beta$ -alanine and carnosine supplementation of mussel

Mytilus coruscus as revealed by UPLC-MS/MS based untargeted metabolomics. Aquaculture Rep. 2021; 20:100730.

<https://doi.org/10.1016/j.aqrep.2021.100730>

**How to Cite This Article:**

Amiri Dashatan N, Ashrafmansouri M, Koushki M, Ahmadi N. Effect of Resveratrol and Its Derivatives on Leishmania Viability: A Meta-Analysis, J Adv Med Biomed Res. 2023; 31(144): 388-394.

**Download citation:**

[BibTeX](#) | [RIS](#) | [EndNote](#) | [Medlars](#) | [ProCite](#) | [Reference Manager](#) | [RefWorks](#)

**Send citation to:**

 [Mendeley](#)  [Zotero](#)  [RefWorks](#)