

Inhibitory Effect of *Syzygium aromaticum* Extract on the Growth of *Pseudomonas aeruginosa* Clinical Isolates Associated with Urinary Tract Infections

Diana Basim Abdulhameed Al-Qaysi^{1*}, Zahraa Qasim Mousa^{2,3}, Zainab Jabbar Sukkar¹

1. Department of Clinical Laboratories Sciences, College of Pharmacy, University of Misan, Maysan, Iraq
2. Department of Biology, College of Science, University of Misan, Maysan, Iraq
3. General Directorate for Education in the province of Maysan, Ministry of Education, Maysan, Iraq



Article Info

doi:10.30699/jambr.33.162.148

Received: 2025/09/26;

Accepted: 2025/11/18;

Published Online: 29 Dec 2025;

Use your device to scan and read the article online



*Corresponding author:

Diana Basim Abdulhameed Al-Qaysi,
Department of Clinical Laboratories
Sciences, College of Pharmacy,
University of Misan, Maysan, Iraq

Email: diana_basim@uomisan.edu.iq

ABSTRACT

Background & Objective: *Syzygium aromaticum* is the scientific name for the aromatic flower buds of an Indonesian native tree from the Myrtaceae family. Cloves contain bioactive compounds such as caryophyllene, eugenol, and acetyl eugenol, which have been shown to possess various physiological properties, including antioxidant, antibacterial, analgesic, anti-inflammatory, anticancer, and anesthetic effects. The present study aims to examine the antibacterial effects of *S. aromaticum* extracts prepared using different solvents (methanol, ethanol, petroleum ether, n-hexane, and chloroform) against *P. aeruginosa*.

Materials & Methods: A total of 93 urine samples were collected from inpatients at Al-Sadder Hospital in Maysan, Iraq. The study specifically targeted urinary tract infections (UTIs) caused by *Pseudomonas aeruginosa*. The antibacterial activity of various *S. aromaticum* extracts (prepared using methanol, ethanol, petroleum ether, hexane, and chloroform) was evaluated against the isolates using the agar well diffusion method. The extracts were tested at concentrations of 62.5, 125, and 500 µg/mL and compared with a standard antibiotic. This investigation aimed to verify the antimicrobial potential of *S. aromaticum* against *P. aeruginosa*, with the findings highlighting its promise for medicinal applications.

Results: Phytochemical screening revealed the presence of tannins, flavonoids, terpenoids, alkaloids, phlobatannins, and saponins in the extracts. Among the tested extracts, the ethanol extract showed the lowest activity against *P. aeruginosa*, producing an inhibition zone of 24 mm at 500 µg/mL. The methanol extract exhibited the highest antibacterial activity, with an inhibition zone of 40 mm at 500 µg/mL. The *P. aeruginosa* isolates were sensitive to Gentamicin, Neomycin, Enrofloxacin, Imipenem, and Meropenem but resistant to Cefadroxil, Cephalexin, Ceftriaxone, Ciprofloxacin, and Sulfamethoxazole.

Conclusion: The results demonstrate notable antibacterial activity of *S. aromaticum* particularly its methanolic extract against *P. aeruginosa*. This potent activity is attributed to its diverse phytochemical composition. Overall, *S. aromaticum* shows potential as a natural source for developing novel antimicrobial agents capable of combating multidrug-resistant infections.

Keywords: *S. aromaticum*, Phytochemical Composition, *Pseudomonas aeruginosa*, Urinary Tract Infections



Copyright © 2025, This is an original open-access article distributed under the terms of the [Creative Commons Attribution-noncommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/) which permits copy and redistribution of the material just in noncommercial usages with proper citation.

1. Introduction

Pseudomonas aeruginosa is a Gram-negative bacterial pathogen responsible for a wide range of diseases, including pneumonia, infections in immunocompromised patients, and complications associated with cystic fibrosis (1). It is also a major cause of high-mortality infections in individuals

with cystic fibrosis, neonatal infections, cancer, and severe burns (2, 3).

During infection, *P. aeruginosa* may adopt either a planktonic lifestyle (typical of acute infections) or a sessile, biofilm-associated lifestyle (common in chronic

infections). Several mechanisms contribute to its virulence, including cytotoxicity (hydrogen cyanide, exotoxin A, T3SS, pyocyanin), immune evasion (elastase, alkaline protease), antibiotic resistance (efflux pumps and modifying enzymes), iron acquisition (siderophores, proteases), biofilm formation (alginate, rhamnolipids), and motility (flagella, pili) (4).

The search for new antibacterial agents—including those derived from medicinal plants—has intensified in response to increasing antimicrobial resistance. Traditional medicine often relies on aromatic herbs to treat bacterial infections, and many plant-based preparations have shown promising therapeutic value (5, 6).

S. aromaticum consists of the dried flower buds of an evergreen tree belonging to the Myrtaceae family, widely grown in tropical regions. For centuries, it has been utilized in traditional medicine and cuisine, and its essential oil is used in perfumery, food flavoring, and herbal treatments. Historically, *S. aromaticum* has been associated with enhancing immunity and promoting resistance to disease. It continues to be used as an anesthetic, antiseptic, antiviral, antifungal, and antibacterial agent for various health conditions (7).

Eugenol, the major volatile component of *S. aromaticum*, is responsible for its characteristic flavor and aroma (8). The flower buds also possess diverse pharmacological properties, including antioxidant, antibacterial, antifungal, anti-inflammatory, anticancer, and antidiabetic activities (9, 10). *S. aromaticum* extracts have shown antibacterial activity against several Gram-negative uropathogens (11).

In addition to tannins, vanillin, flavonoids, and triterpenoids, *S. aromaticum* buds contain 15–20% essential oil, primarily composed of eugenol, eugenol acetate, and caryophyllene. Their distinct aroma is further influenced by trace compounds such as methyl amyl ketone and methyl salicylate (12).

The present study aims to examine the antibacterial effects of *S. aromaticum* extracts prepared using different solvents (methanol, ethanol, petroleum ether, n-hexane, and chloroform) against *P. aeruginosa*. The extracts' efficacy was compared with standard antibiotics, and phytochemical screening was conducted to identify secondary metabolites associated with antibacterial activity. The findings may serve as a basis for further research on identifying, characterizing, and elucidating the mechanisms of active compounds.

2. Materials and Methods

2.1 Preparation of *S. aromaticum*

S. aromaticum buds was purchased from the local market in Maysan, Iraq. The identification and authentication of *S. aromaticum* was confirmed at the Department of Pharmacognosy and Medicinal Plants, College of Pharmacy, University of Misan, Maysan, Iraq.

At first, *S. aromaticum* were carefully washed and cleaned with tap water and then distilled water. Then, they were allowed to air dry. The plant material was finely powdered after drying (13).

2.2 Preparation of test samples and procedure

To identify phytoconstituents, the n-hexane, ethanol, petroleum ether, methanol, and chloroform extracts underwent preliminary phytochemical screening using the following procedure: 50 mL of n-hexane, ethanol, petroleum ether, methanol, and chloroform were added individually as solvents to separate conical flasks containing the test samples (5.0 g each). The mixtures were kept at room temperature for twenty-four hours. The crude clove extracts were then evaporated at 45 °C. After evaporation, the extracted material was stored at 4 °C in the dark. Before antibacterial testing, the extract was dissolved in 10% dimethyl sulfoxide (DMSO) to obtain the required concentrations (14, 15).

2.3 Clinical Bacteria Isolates

Urine samples were collected from patients admitted to the Intensive Care Unit (ICU) and the Burn Unit. The study focused exclusively on samples obtained from patients clinically diagnosed with urinary tract infections (UTIs) caused by *P. aeruginosa*. Samples containing other bacterial pathogens were excluded. The presence of *P. aeruginosa* was confirmed using standard microbiological identification procedures, including culture characteristics, Gram staining, and biochemical testing.

A total of fifteen *P. aeruginosa* isolates were recovered from ninety-three (93) culture-positive urine samples. Of these, eleven (11) isolates exhibiting a multidrug-resistant (MDR) phenotype—as determined by antimicrobial susceptibility testing using the Kirby–Bauer disk diffusion method—were selected for subsequent evaluation of the antibacterial activity of *S. aromaticum* extract.

To ensure the reliability and reproducibility of experimental outcomes, all agar well diffusion assays included a positive control (gentamicin 10 µg discs) and a negative control (Dimethyl Sulfoxide, DMSO). Following inoculation onto blood agar and MacConkey agar, all samples were incubated for 18 hours at 37°C. Plates were then incubated at 37°C for an additional 24 hours. Pure colonies obtained were subcultured onto nutrient agar. Identification of *P. aeruginosa* isolates was confirmed using morphological characteristics, Gram staining, conventional biochemical assays, and the VITEK automated system.

2.4 Biochemical tests

Conventional biochemical tests were performed to confirm the identity of the isolates, including Gram staining, urease production, oxidase test, lactose fermentation, motility test, catalase activity, and microscopic examination (16).

1. To perform the alkaloid test, 0.5 g of extract is weighed and mixed with 1 mL of 2N HCl and 9 mL of distilled water. The mixture is heated in a water bath for two minutes, cooled, and filtered. The filtrate is then used for reactions with Mayer's, Bouchardat's, and Dragendorff's reagents. For Mayer's reagent, three drops of *S. aromaticum* leaf extract and two drops of Mayer's reagent are added to a test tube. The appearance of a white or yellow precipitate indicates the presence of alkaloids. For Bouchardat's reagent, three drops of *S. aromaticum* leaf extract and two drops of the reagent are mixed. A brown to black precipitate indicates alkaloids. For Dragendorff's reagent, three drops of extract and two drops of Dragendorff's reagent are combined. Formation of orange to red or brick-red precipitates confirms the presence of alkaloids. The sample is considered positive for alkaloids if at least two of the three reagents give a positive reaction.

2. For the flavonoid test, 10 mL of hot water is added to a test tube containing the thick *S. aromaticum* extract. The mixture is boiled and filtered while still hot. To 5 mL of filtrate, 0.1 g of Mg powder, 1 mL of HCl, and 2 mL of amyl alcohol are added. After shaking, the mixture is allowed to separate. The presence of flavonoids is indicated by a yellow, orange, or red coloration in the amyl alcohol layer.

3. A thick extract of *S. aromaticum* is placed in a test tube, and hot water is added. The tube is shaken vigorously for 15 minutes. After adding one drop of 2N HCl, persistent foam lasting approximately 10 minutes indicates the presence of saponins.

4. A thick *S. aromaticum* leaf extract is mixed with 10 mL of distilled water and filtered. Distilled water is then added to the filtrate until it becomes colorless. One to two drops of iron (III) chloride reagent are added to 2 mL of the filtrate. A dark blue or blackish-green coloration indicates the presence of tannins.

5. For the triterpenoid test, 30 mg of extract is treated with 5 drops of glacial acetic acid and 2 drops of concentrated sulfuric acid. A blue or green color indicates the presence of steroids, while a red or purple color indicates the presence of triterpenoids.

6. To test for phlobatannins, 500 μ L of the clove plant extract is heated with 0.5 mL of 1% aqueous HCl. The formation of a red precipitate confirms the presence of phlobatannins.

7. For the steroid test, 0.5 g of extract is treated with 2 mL of acetic anhydride, followed by the careful addition of 2 mL of sulfuric acid along the wall of the test tube. A color change from violet to blue-green indicates the presence of steroids.

2.5 Stock and Standard Concentrations of *S. aromaticum* Extract

The stock solution consisted of crude extract at 100 mg/mL⁻¹ DMSO (20mg/50ml) was used to prepare concentrations of 62.5 mg/mL⁻¹, 125 mg/mL, and 500 mg/mL⁻¹ for antibacterial testing.

2.6 Antibacterial Activity of Plant Extracts

The isolates were re-cultured on Mueller–Hinton agar, and antibacterial activity was evaluated using the agar well diffusion method. A 0.5 McFarland suspension (\approx 10⁸ CFU/mL) was prepared and spread on the agar. Wells of 5 mm diameter were filled with 50 μ L of extract. Gentamicin served as the reference antibiotic. Plates were incubated at 35°C \pm 2°C for 18–24 hours and inhibition zones were measured. The germs collected were re-isolated on Muller-Hinton Agar for *Pseudomonas*. The tested extract solutions were prepared. Antibiotic susceptibility testing was performed using the agar well diffusion method described by Karou et al (17). Active extracts were initially identified starting at a high concentration through a presumptive test. A microbial suspension of 0.5 McFarland (\approx 10⁸ CFU/mL) was used. Densitometer turbidity measurements were performed to quantify the suspension. The inoculum was added to culture media prepared according to standard procedures. Sterility and fertility tests were conducted prior to use to assess the quality of the media. Following inoculation of the medium, sterile hollow punches were used to create wells with a diameter of 5 mm. Fifty milliliters of extract at concentrations of 62.5 mg/mL, 125 mg/mL, and 500 mg/mL were added to each well. The reference antibiotic was 10 mg of gentamicin solution for *Pseudomonas*. The Petri dishes were incubated for 18 to 24 hours at 35°C \pm 2°C after 30 minutes of pre-diffusion at room temperature. Diameters of microbial growth inhibition zones were measured using an electronic reading chart.

2.7 Antibiotic resistance

The agar well diffusion method was used to examine the antibacterial activity of five extracts of *C.zeylanicum* bark (18).

Antibiotic susceptibility testing was performed using the agar well diffusion method. Tested antibiotics included Gentamicin (10 mg), Neomycin (30mg), Meropenem (10 mg), Imipenem (10 mg), Cephalexin (30mg), Cefadroxil (30 mg), Ceftriaxone (30 mg), Enrofloxacin (5 mg), Ciprofloxacin (5mg), and Sulfamethoxazole (25 mg) (19, 20).

2.8 Statistical Analysis

All experiments were performed in triplicate (n = 3). Data were expressed as mean \pm SD. One-way ANOVA followed by Tukey's post hoc test was used to determine significant differences (p < 0.05). Statistical analyses were conducted using SPSS version 22.

3. Result

3.1 phytochemical composition of *S. aromaticum*

Based on initial phytochemical analysis, the extract of *S. aromaticum* buds in the current study contain alkaloids, flavonoids, terpenoids, tannins, phlobotannin, steroids, and saponins.

3.2 Sensitivity test for antibiotics of *P. aeruginosa*

The antibiotic susceptibility results of the clinical isolates are summarized in Table 2. Most isolates showed high resistance rates to the tested antibiotics, with only a limited number remaining sensitive. The presence of multidrug-resistant isolates was evident.

3.3 Antibacterial activity of *S. aromaticum* extract against isolates of *P. aeruginosa*

The inhibitory action of *S. aromaticum* extract is determined using three different concentrations of the

extract: 62.5 mg.mL⁻¹, 125 mg.mL⁻¹, and 500 mg.mL⁻¹. Five extracts of Ceylon *S. aromaticum* are used against *P. aeruginosa*, including chloroform, hexane, petroleum ether, methanol, and ethanol. The findings demonstrate that the methanol extract had the largest inhibition range against *P. aeruginosa*, measuring 40 mm at a concentration of 500 mg/ml⁻¹. Petroleum ether came in second with an inhibition range of 33 mm at the same concentration, while the ethanol extract had the smallest inhibition range (Table 3).

Table 1. Phytochemical composition of *S. aromaticum*.

Phytochemical tests	<i>S. aromaticum</i> extracts				
	Ethanol	n-hexane	Methanol	petroleum ether	chloroform
Alkaloids	-	+	+	-	+
Flavonoids	-	-	+	+	+
Saponins	-	+	+	-	+
Steroids	+	-	+	-	+
Tannins	+	+	+	-	-
Phlobotannins	+	-	+	+	-
Terpenoids	-	+	+	-	+

Table 2. Antibiotic sensitivity test of *P. aeruginosa*.

Antibiotics	Clinical Isolate															
	p1	p2	p3	p4	p5	p6	p7	p8	p9	p10	p11	p12	p13	p14	p15	
Cefadroxil (30 mg)	R	R	R	R	R	R	R	R	R	R	R	R	R	R	S	S
Cephalexin (30mg)	R	R	R	R	R	S	R	R	R	R	R	S	S	R	R	
Ceftriaxone (30 mg)	R	R	R	R	R	R	R	R	S	R	R	R	S	S	S	
Ciprofloxacin (5 mg)	R	R	R	R	R	R	S	R	R	R	S	S	R	R	R	
Enrofloxacin (5 mg)	R	R	R	R	R	R	R	R	R	R	R	S	S	S	S	
Sulfamethoxazole (25 mg)	R	R	R	R	R	R	R	R	R	S	R	R	S	S	R	
Neomycin (30 mg)	R	R	R	R	S	R	R	S	S	R	R	S	S	R	S	
Imipinem (10mg)	R	R	R	R	R	S	S	R	R	R	S	S	R	S	S	
Meropenem(10 mg)	R	R	R	R	S	R	R	S	R	R	R	S	S	S	S	

Table 3. Inhibitory effect of *S. aromaticum* extracts.

Concentration (mg/mL)	Ethanol extract	n-Hexane extract	Methanol extract	Petroleum ether extract	Chloroform extract
62.5	15.0 ± 0.5 ^a	15.0 ± 0.4 ^a	31.0 ± 0.6 ^b	27.0 ± 0.5 ^c	27.0 ± 0.5 ^c
125	22.5 ± 0.6 ^a	22.5 ± 0.5 ^a	32.0 ± 0.6 ^b	30.0 ± 0.5 ^b	28.5 ± 0.5 ^b
500	24.0 ± 0.5 ^b	25.5 ± 0.6 ^b	40.0 ± 0.7 ^a	33.0 ± 0.6 ^b	30.0 ± 0.5 ^b

Note: Values are expressed as mean ± SD (n = 3). Different superscript letters within the same row indicate significant differences among extracts at the same concentration (p < 0.05, one-way ANOVA followed by Tukey's test).

4. Discussion

The medicinal value of plants arises from the bioactive chemical compounds they contain, each of which has specific physicochemical effects on the human body Al-Behadili and Faraj (15) and, Batiha et al (9). Phytochemicals exhibit diverse biological activities that may play a role in preventing chronic diseases such as heart disease and stroke. For instance, saponins protect against hypercholesterolemia, while steroids and terpenoids have been shown to exert analgesic effects on the central nervous system. The importance of alkaloids, saponins, and tannins any of which contribute to the action of existing antibiotics—is increasingly recognized by researchers (21).

Flavonoids, a widespread group of phenolic secondary metabolites, are present in the leaves, stems, roots, flowers, and seeds of nearly all plant families. These compounds can exert antibacterial activity by disrupting the cytoplasmic membrane, inhibiting nucleic acid synthesis, and slowing cellular energy metabolism, making them important natural antimicrobial agents (22). The present study demonstrates that *P. aeruginosa* was sensitive to Gentamicin, Neomycin, Meropenem, and Enrofloxacin; however, it exhibited resistance to several other antibiotics tested. *P. aeruginosa* remains a major global health concern due to its ability to resist numerous antibiotics, including imipenem, which is often reserved as a last-line therapeutic option (23).

Given the increasing antibiotic resistance observed in *P. aeruginosa*, the development of new antimicrobial agents and the prudent use of existing antibiotics are essential (9, 24). Findings from this study, along with previous research, indicate that *S. aromaticum* possesses significant antibacterial potential. The results show that the methanolic extract exhibited the greatest inhibitory effect against *P. aeruginosa*, whereas the ethanolic extract displayed the lowest. Previous studies show that *S. aromaticum* extract contains numerous components, predominantly flavonoids and phenolic compounds (25).

Flavonoids play an important role as antimicrobial agents in plants. Serving as part of the plant's natural defense system, they help protect against microbial infections (26). Many plant-derived flavonoids

demonstrate antibacterial effects distinct from those of conventional antibiotics, suggesting their potential use as complementary therapeutic agents (27). Additionally, it is proposed that flavonoid DNA intercalation, which inhibits the synthesis of bacterial nucleic acids, is the mechanism behind the antibacterial properties of flavonoids. The chemical properties of phenolic compounds, notably their aromatic structure and numerous hydroxyl groups capable of donating electrons or hydrogen atoms to neutralize free radicals and other reactive oxygen species (ROS), which are linked to their beneficial bioactivities (15, 28). Furthermore, phenolic compounds exert antibacterial effects by altering virulence factors, genetic regulatory elements, the ability to disrupt biofilms, and the structure and function of membranes (29).

5. Conclusion

Medicinal plants, particularly *S. aromaticum*, hold significant importance in light of the declining effectiveness of many conventional antimicrobial agents. Therefore, exploring new therapeutic alternatives for the treatment of UTIs is essential. The findings of this study highlight the promising antibacterial activity of *S. aromaticum*, especially its methanolic extract, against *P. aeruginosa*. However, isolating and characterizing the active compounds within *S. aromaticum* remains a critical challenge that must be addressed to enable their potential use in biomedical applications.

6. Declarations

6.1 Acknowledgments

The authors would like to express their sincere appreciation to all individuals who contributed to and supported this research.

6.2 Ethical Considerations

The study protocol was approved by the Institutional Review Board (IRB) of University of Misan, Iraq and Ethical code: (No.853-25-5-2025).

6.3 Authors' Contributions

Conceptualization, supervision, funding acquisition and resources: Diana Basim and Zahraa Qasim; Methodology: Diana Basim and Zahraa Qasim and Zainab Jabbar; Data collection: Zainab Jabbar; Data analysis: All authors. All authors reviewed, edited, and approved the final version of the manuscript.

6.4 Conflict of Interest

The authors declare that there is no conflict of interest regarding the publication of this study.

6.5 Fund or Financial Support

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

6.6 Using Artificial Intelligence Tools (AI Tools)

The authors were not utilized AI Tools.

6.7 Availability of Data and Materials

The data that support the findings of this study are available from the corresponding author upon reasonable request.

7. Publisher's Note

This article is part of the Special Issue arising from the Second International Conference for Pharmaceutical Sciences (SICPS 2025), College of Pharmacy, University of Misan, Iraq (29 Nov–1 Dec 2025, see <https://uomisn.edu.iq/pharmacy/conference/>). All manuscripts in this issue were peer-reviewed and accepted for publication in *Journal of Advances in Medical and Biomedical Research (J Adv Med Biomed Res)*.

References

- Reynolds D, Kollef M. The Epidemiology and Pathogenesis and Treatment of Pseudomonas aeruginosa Infections: An Update. *Drugs*. 2021;81(18):2117-31. [PMID] [PMCID] [DOI:10.1007/s40265-021-01635-6]
- Turkina MV, Vikström E. Bacteria-Host Crosstalk: Sensing of the Quorum in the Context of Pseudomonas aeruginosa Infections. *J Innate Immun*. 2019;11(3):263-79. [DOI:10.1159/000494069] [PMID] [PMCID]
- Nathwani D, Raman G, Sulham K, Gavaghan M, Menon V. Clinical and economic consequences of hospital-acquired resistant and multidrug-resistant Pseudomonas aeruginosa infections: a systematic review and meta-analysis. *Antimicrob Resist Infect Control*. 2014;3(1):32. [DOI:10.1186/2047-2994-3-32] [PMID] [PMCID]
- Tam VH, Chang KT, Abdelraouf K, Brioso CG, Ameka M, McCaskey LA, et al. Prevalence, resistance mechanisms, and susceptibility of multidrug-resistant bloodstream isolates of Pseudomonas aeruginosa. *Antimicrob Agents Chemother*. 2010;54(3):1160-4. [PMID] [PMCID] [DOI:10.1128/AAC.01446-09]
- Sannathimmappa MB, Nambiar V, Aravindakshan R. Antibiotics at the crossroads - Do we have any therapeutic alternatives to control the emergence and spread of antimicrobial resistance?. *J Educ Health Promot*. 2021;10:438. [DOI:10.4103/jehp.jehp_557_21]
- Yazarlu O, Iranshahi M, Kashani HRK, Reshadat S, Habtemariam S, Iranshahi M, et al. Perspective on the application of medicinal plants and natural products in wound healing: A mechanistic review. *Pharmacol Res*. 2021;174:105841. [DOI:10.1016/j.phrs.2021.105841] [PMID]
- Vicidomini C, Roviello V, Roviello GN. Molecular Basis of the Therapeutic Potential of Clove (Syzygium aromaticum L.) and Clues to Its Anti-COVID-19 Utility. *Molecules*. 2021;26(7):1880. [PMID] [PMCID] [DOI:10.3390/molecules26071880]
- Ahmed LI, Ibrahim N, Abdel-Salam AB, Fahim KM. Potential application of ginger, clove and thyme essential oils to improve soft cheese microbial safety and sensory characteristics. *Food Biosci*. 2021;42:101177. [DOI:10.1016/j.fbio.2021.101177]
- Batiha GE, Alkazmi LM, Wasef LG, Beshbishy AM, Nadwa EH, Rashwan EK. Syzygium aromaticum L. (Myrtaceae): Traditional Uses, Bioactive Chemical Constituents, Pharmacological and Toxicological Activities. *Biomolecules*. 2020;10(2):202. [PMCID] [DOI:10.3390/biom10020202] [PMID]

10. Maggini V, Semenzato G, Gallo E, Nunziata A, Fani R, Firenzuoli F. Antimicrobial Activity of *Syzygium aromaticum* Essential Oil in Human Health Treatment. *Molecules*. 2024;29(5):999. [DOI:10.3390/molecules29050999] [PMID] [PMCID]
11. Faujdar SS, Bisht D, Sharma A. Antibacterial activity of *Syzygium aromaticum* (clove) against uropathogens producing ESBL, MBL, and AmpC beta-lactamase: Are we close to getting a new antibacterial agent?. *J Family Med Prim Care*. 2020;9(1):180-6. [PMCID] [DOI:10.4103/jfmpc.jfmpc_908_19] [PMID]
12. Cherrat L, Espina L, Bakkali M, Pagán R, Laglaoui A. Chemical composition, antioxidant and antimicrobial properties of *Mentha pulegium*, *Lavandula stoechas* and *Satureja calamintha* Scheele essential oils and an evaluation of their bactericidal effect in combined processes. *Innov Food Sci Emerg Technol*. 2014;22:221-9. [DOI:10.1016/j.ifset.2013.12.016]
13. Rakshit M, Ramalingam C. Screening and comparison of antibacterial activity of Indian spices. *J Exp Sci*. 2010;1(7):33-6.
14. Jasim Al-Tulaibawi NA, Aal-Aaboda M, Abdulhameed Al-Qaesy DB. Prevalence and Antibiotic Sensitivity Profile of Bacteria in Patients with Ear Infections. *J Pure Appl Microbiol*. 2023;17(1):549-53. [DOI:10.22207/JPAM.17.1.52]
15. Al-Behadili WAA, Faraj SH. Phytochemical characterization using GC-MS analysis of bioactive constituents of ethanolic extract of bulbs of *Tulipa gesneriana* L. *Iran J Ichthyol*. 2023;10(Special Issue 1):327-34.
16. Banu KS, Cathrine L. General techniques involved in phytochemical analysis. *Int J Adv Res Chem Sci*. 2015;2(4):25-32.
17. Karou D, Dicko MH, Simporte J, Traore AS. Antioxidant and antibacterial activities of polyphenols from ethnomedicinal plants of Burkina Faso. *Afr J Biotechnol*. 2005;4(8):823-8.
18. Ohikhena FU, Wintola OA, Afolayan AJ. Evaluation of the Antibacterial and Antifungal Properties of *Phragmanthera capitata* (Sprengel) Balle (Loranthaceae), a Mistletoe Growing on Rubber Tree, Using the Dilution Techniques. *Sci World J*. 2017;2017:9658598. [DOI:10.1155/2017/9658598] [PMID] [PMCID]
19. Ebani VV, Nardoni S, Bertelloni F, Najjar B, Pistelli L, Mancianti F. Antibacterial and Antifungal Activity of Essential Oils against Pathogens Responsible for Otitis Externa in Dogs and Cats. *Medicines (Basel)*. 2017;4(2):21. [DOI:10.3390/medicines4020021] [PMID] [PMCID]
20. von Silva-Tarouca MSE, Wolf G, Mueller RS. Determination of minimum inhibitory concentrations for silver sulfadiazine and other topical antimicrobial agents against strains of *Pseudomonas aeruginosa* isolated from canine otitis externa. *Vet Dermatol*. 2019;30(2):145-e42. [DOI:10.1111/vde.12718] [PMID]
21. Erdemoglu N, Ozkan S, Tosun F. Alkaloid profile and antimicrobial activity of *Lupinus angustifolius* L. alkaloid extract. *Phytochem Rev*. 2007;6(1):197-201. [DOI:10.1007/s11101-006-9055-8] [PMCID]
22. Tan Z, Deng J, Ye Q, Zhang Z. The Antibacterial Activity of Natural-derived Flavonoids. *Curr Top Med Chem*. 2022;22(12):1009-19. [PMID] [DOI:10.2174/1568026622666220221110506]
23. Cho JK, Kim JM, Kim KH, Lim HS, Yang CR. Antimicrobial resistance of *Pseudomonas aeruginosa* isolated from dogs and cats. *Korean Soc Vet Serv*. 2021;44(1):21-6.
24. Haro-González JN, Castillo-Herrera GA, Martínez-Velázquez M, Espinosa-Andrews H. Clove Essential Oil (*Syzygium aromaticum* L. Myrtaceae): Extraction, Chemical Composition, Food Applications, and Essential Bioactivity for Human Health. *Molecules*. 2021;26(21):6387. [PMID] [PMCID] [DOI:10.3390/molecules26216387]
25. Al-Tawalbeh DM, Alawneh JM, Momani W, Mayyas A. Comparative antibacterial activity of clove extract against *Pseudomonas aeruginosa*. *BMC Complement Med Ther*. 2025;25(1):7. [PMID] [PMCID] [DOI:10.1186/s12906-024-04740-7]
26. Piasecka A, Jdrzejczak-Rey N, Bednarek P. Secondary metabolites in plant innate immunity: conserved function of divergent chemicals. *New Phytol*. 2015;206(3):948-64. [DOI:10.1111/nph.13325] [PMID]
27. Pandey AK, Shashank K. Perspective on plant products as antimicrobials agents: a review. *Pharmacologia*. 2013;4(7):469-80. [DOI:10.5567/pharmacologia.2013.469.480]
28. Lima MC, Paiva de Sousa C, Fernandez-Prada C, Harel J, Dubreuil JD, de Souza EL. A review of the current evidence of fruit phenolic compounds as potential antimicrobials against pathogenic bacteria. *Microb Pathog*. 2019;130:259-70. [DOI:10.1016/j.micpath.2019.03.025] [PMID]
29. Du W, Zhou M, Liu Z, Chen Y, Li R. Inhibition effects of low concentrations of

epigallocatechin gallate on the biofilm formation and hemolytic activity of *Listeria*

monocytogenes. Food Control. 2018;85:119-26. [[DOI:10.1016/j.foodcont.2017.09.011](https://doi.org/10.1016/j.foodcont.2017.09.011)]

How to Cite This Article:

Al-Qaysi D B A, Mousa Z Q, Sukkar Z J. Inhibitory Effect of *Syzygium aromaticum* Extract on the Growth of *Pseudomonas aeruginosa* Clinical Isolates Associated with Urinary Tract Infections. J Adv Med Biomed Res. 2025;33(162):148-55.

Download citation:

[BibTeX](#) | [RIS](#) | [EndNote](#) | [Medlars](#) | [ProCite](#) | [Reference Manager](#) | [RefWorks](#)

Send citation to:

 [Mendeley](#)  [Zotero](#)  [RefWorks](#) [RefWorks](#)